



Obstructive Sleep Apnea Diagnosis and Treatment

Last Review Date: March 10, 2017

Number: MG.MM.ME.25kv2

Medical Guideline Disclaimer

Property of EmblemHealth. All rights reserved. The treating physician or primary care provider must submit to EmblemHealth the clinical evidence that the patient meets the criteria for the treatment or surgical procedure. Without this documentation and information, EmblemHealth will not be able to properly review the request for prior authorization. The clinical review criteria expressed below reflects how EmblemHealth determines whether certain services or supplies are medically necessary. EmblemHealth established the clinical review criteria based upon a review of currently available clinical information (including clinical outcome studies in the peer-reviewed published medical literature, regulatory status of the technology, evidence-based guidelines of public health and health research agencies, evidence-based guidelines and positions of leading national health professional organizations, views of physicians practicing in relevant clinical areas, and other relevant factors). EmblemHealth expressly reserves the right to revise these conclusions as clinical information changes, and welcomes further relevant information. Each benefit program defines which services are covered. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that this service or supply is covered and/or paid for by EmblemHealth, as some programs exclude coverage for services or supplies that EmblemHealth considers medically necessary. If there is a discrepancy between this guideline and a member's benefits program, the benefits program will govern. In addition, coverage may be mandated by applicable legal requirements of a state, the Federal Government or the Centers for Medicare & Medicaid Services (CMS) for Medicare and Medicaid members. All coding and web site links are accurate at time of publication. EmblemHealth Services Company LLC, ("EmblemHealth") has adopted the herein policy in providing management, administrative and other services to HIP Health Plan of New York, HIP Insurance Company of New York, Group Health Incorporated and GHI HMO Select, related to health benefit plans offered by these entities. All of the aforementioned entities are affiliated companies under common control of EmblemHealth Inc.

Definitions

Apnea	The cessation of airflow for at least 10 seconds.
Hypopnea	An abnormal respiratory event lasting at least 10 seconds associated with at least a 30% reduction in thoracoabdominal movement or airflow as compared to baseline, and with at least a 4% decrease in oxygen saturation.
Apnea-hypopnea index (AHI)	The average number of apneas and hypopneas per hour of sleep without the use of a positive airway pressure device.
Respiratory disturbance index (RDI)	The average number of apneas plus hypopneas per hour of recording without the use of a positive airway pressure device and specifically does NOT include the number of RERAs (respiratory effort related arousals).
Obstructive sleep apnea (OSA)	Characterized by frequent episodes of hypopnea or apnea during sleep. The level of obstruction (retropalatal, retrolingual, nasal or nasopharyngeal) is variable.
Mild apnea	AHI or RDI of 5–14 episodes of apnea or slowed breathing per hour with $\geq 88\%$ oxygen saturation in the blood. Symptoms may include drowsiness or falling asleep during activities that do not require much attention, such as watching TV or reading. These symptoms may cause only minor problems with work or social function.
Moderate apnea	<u>AHI or RDI of 15–30</u> episodes of apnea or slowed breathing per hour with 80% to 85% oxygen saturation in the blood. Symptoms may include drowsiness or falling asleep during activities that require some attention, such as attending a concert or a meeting. These symptoms may cause moderate problems with work or social function.
Severe apnea	AHI or RDI of > 30 episodes of apnea or slowed breathing per hour with $\leq 79\%$ oxygen saturation in the blood. Symptoms may include drowsiness or falling asleep during activities that require active attention, such as eating, talking, driving or walking. These symptoms may cause severe problems with work or social function.

[In-lab sleep facility polysomnography \(PSG\) \(type I\)](#) — technician-attended comprehensive overnight diagnostic sleep test furnished in a sleep laboratory facility. A technologist supervises the recording during sleep time and has the ability to intervene if needed. Type 1 testing includes at least electroencephalography (EEG), electro-oculography (EOG), electromyography (EMG), heart rate or electrocardiography (ECG), airflow, breathing/respiratory effort and arterial oxygen saturation.

[Portable sleep study monitor \(home sleep test \[HST\]\) \(types II, III and IV\)](#) — three categories of portable monitors have been developed for the diagnosis of OSA. HSTs may be technician-attended or unattended.

1. Type II device — monitors and records a minimum of 7 channels (e.g., EEG, EOG, EMG, ECG-heart rate, airflow, respiratory movement/effort and oxygen saturation [SaO2]).
2. Type III device — monitors and records a minimum of 4 channels (e.g., respiratory movement/effort, airflow ECG-heart rate and SaO2).
3. Type IV device — 3 or more channels that allow measurement of AHI/RDI, and must include airflow, respiratory effort and oximetry.

(Note: Type IV devices that do not report AHI/RDI based on direct measurement or airflow or thoracoabdominal movements are excluded unless they are approved by CMS)

Guidelines and [Appendix](#)

(All sleep studies [including attended in-lab facility sleep studies] require prior authorization from EmblemHealth)

Criteria Title	Section	Page	Appendix
OSA diagnostic sleep testing	1	2	Epworth sleepiness scale
Surgical management	2	4	OSA diagnosis
Oral appliance therapy	3	5	OSA treatment
Positive airway pressure (pap) devices Bilevel (BiPAP), demand positive airway pressure (DPAP), variable positive airway pressure (VPAP), adaptive servoventilation (VPAP Adapt SV), auto-titrating positive airway pressure (AutoPap) and Continuous Positive (CPAP)	4	5	Coding
Post OSA treatment surveillance	5	6	N/A

SECTION 1: OSA DIAGNOSTIC SLEEP TESTING

Members are eligible for technician-attended or unattended sleep studies for the diagnosis of OSA when criteria A, B or C ([pediatrics](#)) are met.

A. Unattended (portable monitor) HST — for members ≥ 19 years of age with a high pre-test probability of OSA who do not have atypical or complicating symptoms (1, 2, 3 or 4):

1. Presence of ≥ 3 of the most common symptoms:
 - a. Loud snoring
 - b. Episodes of apnea, choking, gasping, as observed by bed partner
 - c. Excessive daytime fatigue
2. Presence of both:

- a. Loud snoring or witnessed episodes of apnea, choking or gasping
- b. Epworth Scale score ≥ 9 or loud snoring

3. Epworth Scale score > 9 or loud snoring

AND

- a. One of the following:
 - i. Body mass index (BMI) >27
 - ii. Coronary artery disease (angina or myocardial infarction)
 - iii. Cognitive dysfunction
 - iv. Depression
 - v. Diabetes or metabolic syndrome
 - vi. Erectile dysfunction
 - vii. Headaches on awakening
 - viii. Heart failure
 - ix. Hypertension
 - x. Mood disorder
 - xi. Nighttime awakening with gastroesophageal reflux
 - xii. Nocturia
 - xiii. Pulmonary hypertension
 - xiv. Stroke or TIA

4. Presence of both:

- a. Epworth Scale score > 9
- b. Extreme daytime sleepiness

B. Attended (in-lab sleep facility) PSG¹ — for members ≥ 19 years of age with a high pre-test probability of OSA who present with atypical or complicating symptoms. (Criteria “A” must first be met in addition to #s 1 or 2):

- 1. Significant co-morbidities that could degrade accuracy of testing such as either of the following:
 - a. Severe heart failure (EF ≤ 15 or NYHA Class IV)
 - b. Chronic obstructive pulmonary disease and restrictive pulmonary disorders: (FEV1 <30 or PCO₂ > 45)
 - c. Atrial fibrillation
 - d. Significant tachyarrhythmia or bradyarrhythmia
- 2. Symptoms suggestive of sleep related disorders other than OSA; either:
 - a. Abnormal and sudden loss of muscle tone suggestive of narcolepsy
 - b. Limb movements, tonic clonic activity or abnormal sleep behaviors suggesting epilepsy or parasomnia

Cases where unattended monitoring is technically inadequate or fails to establish the diagnosis of OSA in patients with high pretest probability are subject to Medical Director review.

¹ A Split-night PSG, whereby the final portion is utilized for CPAP titration, may be medically necessary. Occasionally, an additional full-night PSG may be necessary for CPAP titration if during the split-night study the vast majority of obstructive respiratory events remained present or if the prescribed CPAP treatment failed to control the member's symptoms.

A video-EEG-PSG (PSG with video monitoring of body positions and extended EEG channels) may be medically necessary to differentiate a diagnosis of paroxysmal arousals or other sleep disruptions that are thought to be seizure related when the initial clinical evaluation and results of a standard EEG are inconclusive.

Limitations/Exclusions

Sleep studies are not medically necessary when snoring or extreme daytime sleepiness are the sole reported symptoms.

C. Attended PSG — for members ≤ 18 years of age when any of the following criteria (1–11) are met:

1. Habitual snoring in association with ≥ 1 of the following (a–e) below:
 - a. Restless or disturbed sleep
 - b. Behavioral disturbance or learning disorders including deterioration in academic performance, attention deficit disorder, hyperactivity disorder
 - c. Frequent awakenings
 - d. Enuresis (bedwetting)
 - e. Growth retardation or failure to thrive
 2. Excessive daytime somnolence or altered mental status not explained by other conditions
 3. Polycythemia not explained by other conditions
 4. Cor pulmonale not explained by other conditions
 5. Witnessed apnea with duration > 2 respiratory cycles
 6. Labored breathing during sleep
 7. Hypertrophy of the tonsils or adenoids in members at significant surgical risk (in order to confirm the presence or absence of OSA) to facilitate clinical management decisions
 8. Suspected congenital central alveolar hypoventilation syndrome or sleep-related hypoventilation due to neuromuscular disease or chest wall deformities
 9. Clinical evidence of a sleep-related breathing disorder in infants who have experienced an apparent life-threatening event
 10. For exclusion of OSA in a member who has undergone adenotonsillectomy for suspected OSA > 8 weeks previously
 11. The initial study was inadequate, equivocal or non-diagnostic and the child's parents or caregiver report that the breathing patterns observed at home were different from those during testing
-

SECTION 2: SURGICAL MANAGEMENT

A. Tonsillectomy (with or without adenoidectomy) for members ≤ 18 years of age and ≥ 1 (See [Tonsillectomy/Adenoidectomy](#) Medical Guideline)

B. Uvulopalatopharyngoplasty (UPPP) — Member must meet all of the following criteria for coverage:

1. Diagnosed OSA
2. One of the following:
 - a. Members with moderate OSA (AHI /RDI 15–30) to severe OSA (AHI/RDI > 30)
 - b. Members with mild OSA (AHI/RDI 5–14) to moderate OSA (AHI/RDI 15–30) with documented symptoms of either:
 - i. Excessive daytime sleepiness, impaired cognition, mood disorders or insomnia.
 - ii. Hypertension, ischemic heart disease or history of stroke
 - c. Failure to respond to or tolerate continuous positive airway pressure (CPAP) or any positive airway pressure (PAP) device, or other appropriate noninvasive treatment
 - d. Counseling from a physician with recognized training in sleep disorders about the potential benefits and risks of the surgery

e. Evidence of retropalatal or combination retropalatal/retrolingual obstruction as the OSA cause. Genioglossal advancement, with or without resuspension of the hyoid bone, may be performed with or instead of UPPP.

Limitations/Exclusions

UPPP for the treatment of snoring in the absence of OSA is not considered medically necessary.

C. Mandibular maxillary osteotomy and advancement —both of the following criteria must be met:

1. Satisfaction of criteria a–d above
2. Evidence of retrolingual obstruction as the OSA cause, or previous failure of UPPP to correct the OSA. Separate repositioning of teeth is not considered necessary except under unusual circumstances, but is covered if necessary. Additionally, application of an interdental fixation device is occasionally necessary and is a covered service (see Section 2: [Oral Appliance Therapy](#))

D. Tracheostomy — may be indicated for OSA if, in the judgment of the attending physician, the member is unresponsive to other means of treatment, or in cases where other means of treatment would be ineffective or contraindicated.

When OSA is caused by discrete anatomic abnormalities of the upper airway (e.g., enlarged tonsils or enlarged tongue), surgery to correct these abnormalities is covered if medically necessary, based on adequate documentation in the medical record supporting the significant contribution.

Limitations/Exclusions

The following procedures are not considered medically necessary, as they are regarded as investigational:

1. Palatal implant or stiffening procedures
2. Electro-sleep therapy
3. Laser-assisted uvulopalatoplasty
4. Radiofrequency tissue-volume reduction somnoplasty for upper airway obstruction

Covered for **Medicaid members only** when **one** of the following criteria is met;:

- a. Members with moderate OSA (AHI/RDI 15–30) to severe OSA (AHI/RDI > 30)
 - b. Members with mild OSA (AHI/RDI 5–14) to moderate OSA (AHI/RDI 15–30) with documented symptoms of either:
 - i. Excessive daytime sleepiness, impaired cognition, mood disorders or insomnia.
 - ii. Hypertension, ischemic heart disease or history of stroke
 - c. Failure to respond to or tolerate continuous positive airway pressure (CPAP) or any positive airway pressure (PAP) device, or other appropriate noninvasive treatment
 - d. Counseling from a physician with recognized training in sleep disorders about the potential benefits and risks of the surgery
 - e. Evidence of retrolingual obstruction as the OSA cause
5. Tongue suspension/suturing procedures
 6. Hypoglossal stimulators.

SECTION 3: ORAL APPLIANCE THERAPY

Members are eligible for custom-fitted oral appliances for OSA for either of the following indications:

1. Members with mild asymptomatic OSA (AHI/RDI 5–14; see Section # 4 — CPAP, BiPAP)
2. Members with moderate OSA (AHI/RDI 15–30) to severe OSA (AHI/RDI > 30) who have had a trial of nasal CPAP or any PAP device but are intolerant to treatment

Oral appliance therapy is also indicated for members who are not candidates for tonsillectomy and adenoidectomy, craniofacial operations or tracheostomy.

Limitations/Exclusions

Oral appliance therapy for members with primary snoring (characterized by loud upper-airway breathing sounds in sleep without episodes of apnea) is not considered medically necessary.

SECTION 4: POSITIVE AIRWAY PRESSURE (PAP) DEVICES — Bilevel (BiPAP), demand positive airway pressure (DPAP), variable positive airway pressure (VPAP), adaptive servoventilation (VPAP Adapt SV), auto-titrating positive airway pressure (AutoPAP) and Continuous Positive (CPAP)

Members with the DME benefit are eligible for PAP device coverage when the following criteria are applicable.

1. CPAP: Positive OSA diagnosis and either:
 - a. Members with moderate–severe OSA (AHI/RDI ≥ 15)
 - b. Members with mild OSA (AHI/RDI 5–14) with documented symptoms of either:
 - i. Excessive daytime sleepiness, impaired cognition, mood disorders or insomnia
 - ii. Hypertension, ischemic heart disease or history of stroke

Either an unheated or heated humidifier is covered when ordered by the treating physician for use with a covered PAP device.

2. BiPAP (or similar device): Can be used for sleep apnea instead of CPAP under either of the following circumstances:
 - a. Documentation of failure to eliminate OSA with CPAP pressure of < 20 cm H₂O
 - b. Failure to tolerate CPAP after both a clinical trial and a CPAP titration study

Continued CPAP Coverage — Conversion from rental to purchase

CPAP compliance is defined as use of CPAP ≥ 4 hours per night on 70% of nights during a consecutive thirty (30) day period anytime during the first three (3) months of initial usage.

Adherence to therapy is evidenced by a CPAP Compliance Report detailing hours of usage per night based on actual nights used.

A CPAP device will be purchased if adherence to therapy within the (90) day period is demonstrated per the report. Failure to achieve compliance within this period will result in the denial of the device as not medically necessary.

Members should receive a face-to-face clinical re-evaluation by the treating physician within 2 months of initiating therapy.

Accessories

A7032/A7033	2 per 1 month
A7034	1 per 3 months
A7035	1 per 6 months
A7036	1 per 6 months
A7037	1 per 3months

A7038	2 per 1 month	CPAP accessories are covered when the coverage criteria for the device are met.
A7039	1 per 6 months	The table represents the usual maximum amount of accessories expected to be medically necessary.

Limitations/Exclusions

1. A CPAP device that is obtained if the criteria have not been met will be denied as not medically necessary.
2. Accessories used with the CPAP device will be denied as not medically necessary if they are obtained when the CPAP criteria have not been met.
3. Quantities of supplies greater than those described in this guideline as the usual maximum amounts, in the absence of documentation clearly explaining the medical necessity of the excess quantities, will be denied as not medically necessary.

SECTION 5: POST OSA TREATMENT SURVEILLANCE

Repeat sleep studies may be considered medically necessary up to two times a year when any of the following are applicable:

1. To evaluate PAP treatment effectiveness
2. To determine whether PAP treatment settings require adjustment
3. To determine whether PAP treatment continuation is necessary
4. To assess treatment response post upper airway surgical procedures and after initial treatment with oral appliances

Diagnostic and Procedure Coding

Revision History

3/17/2017 — added pediatric PSG criteria.

8/12/2016 — added coverage criteria for radiofrequency tissue-volume reduction somnoplasty (Medicaid members only)

3/11/2016 — added tonsillectomy (with or without adenoidectomy) to surgical management section.

10/9/2015 — added atrial fibrillation and significant tachyarrhythmia or bradyarrhythmia to list of co-morbidities that could degrade accuracy of testing in the attended sleep lab environment.

9/11/2015 — added hypoglossal nerve stimulators to list of investigational procedures.

References

American Academy of Sleep Medicine. Clinical Guideline for the Use of Unattended Portable Monitors in the Diagnosis of Obstructive Sleep Apnea in Adult Patients. 2007. <http://www.aasmnet.org/jcsm/AcceptedPapers/PMProof.pdf>. Accessed March 22, 2017.

American Academy of Sleep Medicine. An Overview of the Revised Portable Monitoring (PM) Device Practice Parameters - American Academy of Sleep Medicine Recommendations. 2008. <http://www.aastweb.org/Resources/FocusGroups/OverviewRevisedPMPracParams.pdf>. Accessed March 22, 2017.

Canadian Agency for Drugs and Technologies in Health (CADTH). Portable Monitoring Devices for Diagnosis of Obstructive Sleep Apnea at Home: Review of Accuracy, Cost-Effectiveness, Guidelines, and Coverage in Canada. December 2009. <https://cadth.ca/portable-monitoring-devices-diagnosis-obstructive-sleep-apnea-home-review-accuracy-cost-0>. Accessed March 21, 2012.

CMS Decision Memo for Sleep Testing for Obstructive Sleep Apnea (OSA) (CAG-00405N). 2009. <http://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=330&ncdver=1&bc=AgAAgAAAAAAAAA%3d%3d&>. Accessed March 22, 2017.

Bailey BJ, Calhoun KH, Healy GB, et al, eds. *Head and Neck Surgery—Otolaryngology*. Vol 1. 3rd ed. Philadelphia: Lippincott Williams & Wilkins; 2001:594-597.

Balk EM, Moorthy D, Obadan NO, et al. Diagnosis and Treatment of Obstructive Sleep Apnea in Adults. Comparative Effectiveness Review No. 32. Prepared by Tufts Evidence-based Practice Center under Contract No. 290-2007-10055-1. AHRQ Publication No. 11-EHC052-EF. Rockville, MD: Agency for Healthcare Research and Quality; July 2011.

Collop NA, Anderson WM, et al. Clinical Guidelines for the use of unattended portable monitors in the Diagnosis of Obstructive Sleep Apnea in Adult Patients. *J Clin Sleep Med*. 2007(7): 737- 47.

Davila DG. Medical considerations in surgery for sleep apnea. *Atlas Oral Maxillofac Surg Clin North Am*. 1995;7:205-221.

Flemons WW, Littner MR, Rowley JA, et al. Home diagnosis of sleep apnea: a systematic review of the literature. An evidence review cosponsored by the American Academy of Sleep Medicine, the American College of Chest Physicians, and the American Thoracic Society. *Chest* 2003; 124:1543-1579

Kushida CA, Littner MR, Morgenthaler T, et al. Practice Parameters for the Indications of Polysomnography and Related Procedures: an update for 2005. *sleep*; 28: 499-521.

Loube D. Radiofrequency ablation for sleep-disordered breathing. *Chest*. 1998;113:1151-1152.

Marcus CL, Brooks LJ, Draper KA, et al.; American Academy of Pediatrics. Diagnosis and Management of Childhood Obstructive Sleep Apnea Syndrome. *Pediatrics*. 2012;130(3):576-584.

Mulgrew AT, Fox N, Ayas NT, Ryan CF. Diagnosis and Treatment of Obstructive Sleep Apnea without Polysomnography. *Ann Intern Med* 2007; 146: 157-166.

NHIC. LCD for Positive Airway Pressure (PAP) devices for the Treatment of Obstructive Sleep Apnea. October 2014. <http://www.medicarenhic.com/dme/mrlcdcurrent.aspx>. Accessed March 22, 2017.

Roland PS, Rosenfeld RM, Brooks LJ, et al; American Academy of Otolaryngology—Head and Neck Surgery Foundation. Clinical practice guideline: polysomnography for sleep-disordered breathing prior to tonsillectomy in children. *Otolaryngol Head Neck Surg*. 2011 Jul;145(1 Suppl):S1-15.

Sher AE, Schechtman KB, Piccirillo JF. The efficacy of surgical modifications of the upper airway in adults with obstructive sleep apnea syndrome. *Sleep*. 1996;19:156-177.

Sleep-related breathing disorders in adults: recommendations for syndrome definition and measurement techniques in clinical research. The Report of an American Academy of Sleep Medicine Task Force. *Sleep*. 1999;22:667–689.

Specialty-matched clinical peer review.

Trikalinos TA, Ip S, et al. Technology Assessment: Home Diagnosis of Obstructive Sleep Apnea-Hypopnea Syndrome. Agency Healthcare Research Quality. August 2007.

APPENDIX

[Epworth Sleep Scale](#): Page 10

Clinical Pathway: OSA Diagnosis: Page 11

Clinical Pathway: OSA Treatment: Page 12

[Coding](#): Page 12

Epworth Sleepiness Scale

Use the following scale to choose the most appropriate number for each situation:

0 = would **never** doze or sleep


1= **slight** chance of dozing or sleeping

2 = **moderate** chance of dozing or sleeping

3 = **high** chance of dozing or sleeping

Situation	Chance of sleeping or dozing
Sitting and reading	
Watching TV	
Sitting inactive in a public place	
Being a passenger in a motor vehicle for an hour or more	
Lying down in the afternoon	
Sitting and talking to someone	
Sitting quietly after lunch (no alcohol)	
Stopped for a few minutes in traffic while driving	
Total the points for your Epworth Scale	Epworth Score =

Part I: Diagnosis of Obstructive Sleep Apnea




Notes

*There is consensus that sleep testing is appropriate in patients who have signs and symptoms suggestive of OSA. Sleep testing is not appropriate for general population screening or for all patients who snore. There has not been universal agreement as to a minimal set of signs and symptoms that should trigger a sleep study.

The clusters of signs, symptoms, and associated conditions as listed on these pages represent typical and reasonable scenarios. Individual cases may require exceptions to this algorithm.

**The Epworth Sleepiness Scale is a tool to assist clinicians in quantifying sleepiness which can be useful for assessing the need for OSA testing and the response to OSA interventions. It indicates the likelihood of falling asleep in commonly encountered situations. Another tool, the Berlin questionnaire includes questions regarding weight change, snoring loudness, snoring frequency, witnessed breathing pauses during sleep, day time fatigue and hypertension. These responses correlate with subsequently demonstrated OSA on sleep testing

Part II: Treatment of Obstructive Sleep Apnea



¹CPAP titration study may be indicated
²Indications for dental device
 1. Milder OSA
 2. Positional Apnea
 3. Absence of Nasal Obstruction

Applicable Procedure Codes

21121	Genioplasty; sliding osteotomy, single piece
21122	Genioplasty; sliding osteotomies, 2 or more osteotomies (eg, wedge excision or bone wedge reversal for asymmetrical chin)
21141	Reconstruction midface, LeFort I; single piece, segment movement in any direction (eg, for Long Face Syndrome), without bone graft
21196	Reconstruction of mandibular rami and/or body, sagittal split; with internal rigid fixation
21198	Osteotomy, mandible, segmental
21199	Osteotomy, mandible, segmental; with genioglossus advancement
31600	Tracheostomy, planned (separate procedure);
31601	Tracheostomy, planned (separate procedure); younger than two years
42145	Palatopharyngoplasty (eg, uvulopalatopharyngoplasty, uvulopharyngoplasty)
42820	Tonsillectomy and adenoidectomy; younger than age 12
42821	Tonsillectomy and adenoidectomy; age 12 or over ¹
42825	Tonsillectomy, primary or secondary; younger than age 12
42826	Tonsillectomy, primary or secondary; age 12 or over
94660	Continuous positive airway pressure ventilation (CPAP), initiation and management
95782	Polysomnography; younger than 6 years, sleep staging with 4 or more additional parameters of sleep, attended by a technologist Note: Service not included in Medicaid managed care benefit package package prior to 01-01-2016. Services covered for Medicaid as of 01/01/2016
95783	Polysomnography; younger than 6 years, sleep staging with 4 or more additional parameters of sleep, with initiation of continuous positive airway pressure therapy or bi-level ventilation, attended by a technologist Note: Service not included in Medicaid managed care benefit package package prior to 01-01-2016. Services covered for Medicaid as of 01/01/2016
95800	Sleep study, unattended, simultaneous recording; heart rate, oxygen saturation, respiratory analysis (eg, by airflow or peripheral arterial tone), and sleep time
95801	Sleep study, unattended, simultaneous recording; minimum of heart rate, oxygen saturation, and respiratory analysis (eg, by airflow or peripheral arterial tone)
95805	Multiple sleep latency or maintenance of wakefulness testing, recording, analysis and interpretation of physiological measurements of sleep during multiple trials to assess sleepiness Multiple sleep latency or maintenance of wakefulness testing, recording, analysis and interpretation of physiological measurements of sleep during multiple trials to assess sleepiness
95806	Sleep study, unattended, simultaneous recording of, heart rate, oxygen saturation, respiratory airflow, and respiratory effort (eg, thoracoabdominal movement)
95807	Sleep study, simultaneous recording of ventilation, respiratory effort, ECG or heart rate, and oxygen saturation, attended by a technologist
95808	Polysomnography; sleep staging with 1-3 additional parameters of sleep, attended by a technologist
95810	Polysomnography; sleep staging with 4 or more additional parameters of sleep, attended by a technologist
95811	Polysomnography; sleep staging with 4 or more additional parameters of sleep, with initiation of continuous positive airway pressure therapy or bilevel ventilation, attended by a technologist

A4604	Tubing with integrated heating element for use with positive airway pressure device
A7027	Combination oral/nasal mask, used with continuous positive airway pressure device, each
A7028	Oral cushion for combination oral/nasal mask, replacement only, each
A7029	Nasal pillows for combination oral/nasal mask, replacement only, pair
A7030	Full face mask used with positive airway pressure device, each
A7031	Face mask interface, replacement for full face mask, each
A7032	Cushion for use on nasal mask interface, replacement only, each
A7033	Pillow for use on nasal cannula type interface, replacement only, pair
A7034	Nasal interface (mask or cannula type) used with positive airway pressure device, with or without head strap
A7035	Headgear used with positive airway pressure device
A7036	Chinstrap used with positive airway pressure device
A7037	Tubing used with positive airway pressure device
A7038	Filter, disposable, used with positive airway pressure device
A7039	Filter, non disposable, used with positive airway pressure device
A7044	Oral interface used with positive airway pressure device, each
A7045	Exhalation port with or without swivel used with accessories for positive airway devices, replacement only
A7046	Water chamber for humidifier, used with positive airway pressure device, replacement, each
E0470	Respiratory assist device, bi-level pressure capability, without backup rate feature, used with noninvasive interface, e.g., nasal or facial mask (intermittent assist device with continuous positive airway pressure device)
E0471	Respiratory assist device, bi-level pressure capability, with back-up rate feature, used with noninvasive interface, e.g., nasal or facial mask (intermittent assist device with continuous positive airway pressure device)
E0472	Respiratory assist device, bi-level pressure capability, with backup rate feature, used with invasive interface, e.g., tracheostomy tube (intermittent assist device with continuous positive airway pressure device)
E0485	Oral device/appliance used to reduce upper airway collapsibility, adjustable or non-adjustable, prefabricated, includes fitting and adjustment
E0486	Oral device/appliance used to reduce upper airway collapsibility, adjustable or non-adjustable, custom fabricated, includes fitting and adjustment
E0561	Humidifier, non heated, used with positive airway pressure device
E0562	Humidifier, heated, used with positive airway pressure device
E0601	Continuous airway pressure (CPAP) device
G0398	Home sleep study test (HST) with type II portable monitor, unattended; minimum of 7 channels: EEG, EOG, EMG, ECG/heart rate, airflow, respiratory effort and oxygen saturation
G0399	Home sleep test (HST) with type III portable monitor, unattended; minimum of 4 channels: 2 respiratory movement/airflow, 1 ECG/heart rate and 1 oxygen saturation
G0400	Home sleep test (HST) with type IV portable monitor, unattended; minimum of 3 channels

Applicable Diagnosis Codes

E66.01	Morbid (severe) obesity due to excess calories
E66.2	Morbid (severe) obesity with alveolar hypoventilation
F10.182	Alcohol abuse with alcohol-induced sleep disorder
F10.282	Alcohol dependence with alcohol-induced sleep disorder
F10.982	Alcohol use, unspecified with alcohol-induced sleep disorder
F11.182	Opioid abuse with opioid-induced sleep disorder
F11.282	Opioid dependence with opioid-induced sleep disorder
F11.982	Opioid use, unspecified with opioid-induced sleep disorder
F13.182	Sedative, hypnotic or anxiolytic abuse with sedative, hypnotic or anxiolytic-induced sleep disorder
F13.282	Sedative, hypnotic or anxiolytic dependence with sedative, hypnotic or anxiolytic-induced sleep disorder
F13.982	Sedative, hypnotic or anxiolytic use, unspecified with sedative, hypnotic or anxiolytic-induced sleep disorder
F14.182	Cocaine abuse with cocaine-induced sleep disorder
F14.282	Cocaine dependence with cocaine-induced sleep disorder
F14.982	Cocaine use, unspecified with cocaine-induced sleep disorder
F15.182	Other stimulant abuse with stimulant-induced sleep disorder
F15.282	Other stimulant dependence with stimulant-induced sleep disorder
F15.982	Other stimulant use, unspecified with stimulant-induced sleep disorder
F19.182	Other psychoactive substance abuse with psychoactive substance-induced sleep disorder
F19.282	Other psychoactive substance dependence with psychoactive substance-induced sleep disorder
F19.982	Other psychoactive substance use, unspecified with psychoactive substance-induced sleep disorder
F51	Sleep disorders not due to a substance or known physiological condition
F51.0	Insomnia not due to a substance or known physiological condition
F51.01	Primary insomnia
F51.02	Adjustment insomnia
F51.03	Paradoxical insomnia
F51.04	Psychophysilogic insomnia
F51.05	Insomnia due to other mental disorder
F51.09	Other insomnia not due to a substance or known physiological condition
F51.1	Hypersomnia not due to a substance or known physiological condition
F51.11	Primary hypersomnia
F51.12	Insufficient sleep syndrome
F51.13	Hypersomnia due to other mental disorder
F51.19	Other hypersomnia not due to a substance or known physiological condition
F51.3	Sleepwalking [somnambulism]
F51.4	Sleep terrors [night terrors]
F51.5	Nightmare disorder
F51.8	Other sleep disorders not due to a substance or known physiological condition
F51.9	Sleep disorder not due to a substance or known physiological condition, unspecified
G12.0	Infantile spinal muscular atrophy, type I [Werdnig-Hoffman]

G12.1	Other inherited spinal muscular atrophy
G12.21	Amyotrophic lateral sclerosis
G12.22	Progressive bulbar palsy
G12.29	Other motor neuron disease
G12.8	Other spinal muscular atrophies and related syndromes
G25.3	Myoclonus
G25.81	Restless legs syndrome
G37.3	Acute transverse myelitis in demyelinating disease of central nervous system
G47.00	Insomnia, unspecified
G47.01	Insomnia due to medical condition
G47.09	Other insomnia
G47.1	Hypersomnia
G47.10	Hypersomnia, unspecified
G47.11	Idiopathic hypersomnia with long sleep time
G47.12	Idiopathic hypersomnia without long sleep time
G47.13	Recurrent hypersomnia
G47.14	Hypersomnia due to medical condition
G47.19	Other hypersomnia
G47.2	Circadian rhythm sleep disorders
G47.20	Circadian rhythm sleep disorder, unspecified type
G47.21	Circadian rhythm sleep disorder, delayed sleep phase type
G47.22	Circadian rhythm sleep disorder, advanced sleep phase type
G47.23	Circadian rhythm sleep disorder, irregular sleep wake type
G47.24	Circadian rhythm sleep disorder, free running type
G47.25	Circadian rhythm sleep disorder, jet lag type
G47.26	Circadian rhythm sleep disorder, shift work type
G47.27	Circadian rhythm sleep disorder in conditions classified elsewhere
G47.29	Other circadian rhythm sleep disorder
G47.3	Sleep apnea
G47.30	Sleep apnea, unspecified
G47.31	Primary central sleep apnea
G47.32	High altitude periodic breathing
G47.33	Obstructive sleep apnea (adult) (pediatric)
G47.34	Idiopathic sleep related nonobstructive alveolar hypoventilation
G47.35	Congenital central alveolar hypoventilation syndrome
G47.36	Sleep related hypoventilation in conditions classified elsewhere
G47.37	Central sleep apnea in conditions classified elsewhere
G47.39	Other sleep apnea
G47.4	Narcolepsy and cataplexy
G47.41	Narcolepsy
G47.411	Narcolepsy with cataplexy

G47.419	Narcolepsy without cataplexy
G47.42	Narcolepsy in conditions classified elsewhere
G47.421	Narcolepsy in conditions classified elsewhere with cataplexy
G47.429	Narcolepsy in conditions classified elsewhere without cataplexy
G47.5	Parasomnia
G47.50	Parasomnia, unspecified
G47.51	Confusional arousals
G47.52	REM sleep behavior disorder
G47.53	Recurrent isolated sleep paralysis
G47.54	Parasomnia in conditions classified elsewhere
G47.59	Other parasomnia
G47.6	Sleep related movement disorders
G47.61	Periodic limb movement disorder
G47.62	Sleep related leg cramps
G47.63	Sleep related bruxism
G47.69	Other sleep related movement disorders
G47.8	Other sleep disorders
G47.9	Sleep disorder, unspecified
G71.0	Muscular dystrophy
G71.11	Myotonic muscular dystrophy
G71.2	Congenital myopathies
G71.9	Primary disorder of muscle, unspecified
G72.9	Myopathy, unspecified
G80.9	Cerebral palsy, unspecified
G82.50	Quadriplegia, unspecified
G90.1	Familial dysautonomia [Riley-Day]
G93.1	Anoxic brain damage, not elsewhere classified
I42.0	Dilated cardiomyopathy
I42.5	Other restrictive cardiomyopathy
I42.7	Cardiomyopathy due to drug and external agent
I42.8	Other cardiomyopathies
I42.9	Cardiomyopathy, unspecified
I50.2	Systolic (congestive) heart failure
I50.20	Unspecified systolic (congestive) heart failure
I50.21	Acute systolic (congestive) heart failure
I50.22	Chronic systolic (congestive) heart failure
I50.23	Acute on chronic systolic (congestive) heart failure
I50.3	Diastolic (congestive) heart failure
I50.30	Unspecified diastolic (congestive) heart failure
I50.31	Acute diastolic (congestive) heart failure
I50.32	Chronic diastolic (congestive) heart failure

I50.33	Acute on chronic diastolic (congestive) heart failure
I50.4	Combined systolic (congestive) and diastolic (congestive) heart failure
I50.40	Unspecified combined systolic (congestive) and diastolic (congestive) heart failure
I50.41	Acute combined systolic (congestive) and diastolic (congestive) heart failure
I50.42	Chronic combined systolic (congestive) and diastolic (congestive) heart failure
I50.43	Acute on chronic combined systolic (congestive) and diastolic (congestive) heart failure
I50.9	Heart failure, unspecified
J35.3	Hypertrophy of tonsils with hypertrophy of adenoids
J96.1	Chronic respiratory failure
J96.10	Chronic respiratory failure, unspecified whether with hypoxia or hypercapnia
J96.11	Chronic respiratory failure with hypoxia
J96.12	Chronic respiratory failure with hypercapnia
N52	Male erectile dysfunction
N52.0	Vasculogenic erectile dysfunction
N52.01	Erectile dysfunction due to arterial insufficiency
N52.02	Corporo-venous occlusive erectile dysfunction
N52.03	Combined arterial insufficiency and corporo-venous occlusive erectile dysfunction
N52.1	Erectile dysfunction due to diseases classified elsewhere
N52.2	Drug-induced erectile dysfunction
N52.3	Post-surgical erectile dysfunction
N52.31	Erectile dysfunction following radical prostatectomy
N52.32	Erectile dysfunction following radical cystectomy
N52.33	Erectile dysfunction following urethral surgery
N52.34	Erectile dysfunction following simple prostatectomy
N52.39	Other post-surgical erectile dysfunction
N52.8	Other male erectile dysfunction
N52.9	Male erectile dysfunction, unspecified
Q07.8	Other specified congenital malformations of nervous system
Q31.1	Congenital subglottic stenosis
Q31.2	Laryngeal hypoplasia
Q31.3	Laryngocele
Q31.5	Congenital laryngomalacia
Q31.8	Other congenital malformations of larynx
Q31.9	Congenital malformation of larynx, unspecified
Q32	Congenital malformations of trachea and bronchus
Q32.0	Congenital tracheomalacia
Q32.1	Other congenital malformations of trachea
Q32.2	Congenital bronchomalacia
Q32.3	Congenital stenosis of bronchus
Q32.4	Other congenital malformations of bronchus
Q75	Other congenital malformations of skull and face bones

Q75.0	Craniosynostosis
Q75.1	Craniofacial dysostosis
Q75.2	Hypertelorism
Q75.3	Macrocephaly
Q75.4	Mandibulofacial dysostosis
Q75.5	Oculomandibular dysostosis
Q75.8	Other specified congenital malformations of skull and face bones
Q75.9	Congenital malformation of skull and face bones, unspecified
Q77.0	Achondrogenesis
Q77.1	Thanatophoric short stature
Q77.4	Achondroplasia
Q77.5	Diastrophic dysplasia
Q77.7	Spondyloepiphyseal dysplasia
Q77.8	Other osteochondrodysplasia with defects of growth of tubular bones and spine
Q77.9	Osteochondrodysplasia with defects of growth of tubular bones and spine, unspecified
Q78.4	Enchondromatosis
Q78.9	Osteochondrodysplasia, unspecified
Q87.0	Congenital malformation syndromes predominantly affecting facial appearance
R06.00	Dyspnea, unspecified
R06.09	Other forms of dyspnea
R06.3	Periodic breathing
R06.81	Apnea, not elsewhere classified
R06.83	Snoring
R06.89	Other abnormalities of breathing
R09.02	Hypoxemia