

Septoplasty

Last Review Date: August 11, 2017

Number: MG.MM.SU.04dC

Medical Guideline Disclaimer

Property of EmblemHealth. All rights reserved. The treating physician or primary care provider must submit to EmblemHealth the clinical evidence that the patient meets the criteria for the treatment or surgical procedure. Without this documentation and information, EmblemHealth will not be able to properly review the request for prior authorization. The clinical review criteria expressed below reflects how EmblemHealth determines whether certain services or supplies are medically necessary. EmblemHealth established the clinical review criteria based upon a review of currently available clinical information (including clinical outcome studies in the peer-reviewed published medical literature, regulatory status of the technology, evidence-based guidelines of public health and health research agencies, evidence-based guidelines and positions of leading national health professional organizations, views of physicians practicing in relevant clinical areas, and other relevant factors). EmblemHealth expressly reserves the right to revise these conclusions as clinical information changes, and welcomes further relevant information. Each benefit program defines which services are covered. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that this service or supply is covered and/or paid for by EmblemHealth, as some programs exclude coverage for services or supplies that EmblemHealth considers medically necessary. If there is a discrepancy between this guideline and a member's benefits program, the benefits program will govern. In addition, coverage may be mandated by applicable legal requirements of a state, the Federal Government or the Centers for Medicare & Medicaid Services (CMS) for Medicare and Medicaid members. All coding and web site links are accurate at time of publication. EmblemHealth Services Company LLC, ("EmblemHealth") has adopted the herein policy in providing management, administrative and other services to HIP Health Plan of New York, HIP Insurance Company of New York, Group Health Incorporated and GHI HMO Select, related to health benefit plans offered by these entities. All of the aforementioned entities are affiliated companies under common control of EmblemHealth Inc.

Definition

Septoplasty (submucosal resection of the septum) is the surgical correction or reconstruction of the nasal septum, which divides the right and left cavities. The septum is generally situated in the direct center of the nose; however, when the septum is off-center or misaligned, corrective surgery is sometimes required to correct the breathing impairment that results from the misalignment.

Related Guidelines

[Cosmetic Surgery Procedures](#)

[Rhinoplasty](#)

Guideline

Members are eligible for coverage of septoplasty when **at least 1** of the following conditions exists:

- I. Reconstruction post tumor or polyp removal or surgical excision of part of the ethmoid bone (ethmoidectomy or turbinate reduction).
- II. Chronic nasal obstruction attributable to deviated septum that results in significant medical disabilities, such as:
 1. **Recurrent purulent sinusitis**¹ (> 3 episodes per year) with > 50% middle meatus complex obstruction on the same side. Conservative management has failed to alleviate or prevent episodes of sinusitis, including **all**:
 - Antibiotics

¹ Diagnosis must be documented by both of the following: Symptoms (including purulent nasal drainage) and radiologic evidence of either recurrent acute sinusitis (> 3 episodes per year) or chronic rhinosinusitis (symptoms > 12 weeks (clouding of sinuses, thickening of sinus membranes on limited CT).

- Antihistamines, decongestants, nasal sprays or topical steroids
 - Attempts to discontinue nasal irritants such as smoking, drugs and inadequate humidification
2. **Nasal septal deformity or spur (congenital or acquired) with > 50% airway obstruction; all:**
 - Assessment and testing, if appropriate, for allergic conditions completed
 - Conservative management failure in the presence of allergic rhinitis (i.e., allergic precautions, antihistamines, topical nasal steroids and plus/minus desensitization injections)
 - General conservative management failure (i.e., reduction of all nasal irritants such as smoking, occupational exposures, drugs and inadequate humidification)
 3. **Recurrent Epistaxis (at least 4, or an episode of severe epistaxis associated with a septal deviation or spur seen on exam); both:**
 - Conservative treatment measure failure, such as avoidance of medications affecting coagulation, addition of environmental humidity and cauterization
 - Nasal spur or septal deformity etiology of abnormal airflow
 4. **Unusual facial pain** originating from the nasal area and relieved by septal anesthesia.
 5. **Impending septal perforation (hole through the septum); both:**
 - Significant septal deviation with airflow obstruction
 - Conservative measure failure, including humidification, desisting further digital trauma, reduction of nasal irritants (smoking, drugs) and drug therapy (decongestants, antihistamines, nasal steroids), if indicated
- III. Asymptomatic deformity that prevents surgical access to other intranasal or paranasal areas (e.g., sinuses, turbinates)
- IV. When performed in association with cleft palate repair

Limitations/Exclusions

Septoplasty is not considered medically necessary for any indications other than those listed above.

Revision History

8/12/2016: Added septoplasty coverage when performed with cleft palate repair, as well as when an asymptomatic deformity impacts surgical access to other intranasal/paranasal areas.

Applicable Procedure Codes

30520	Septoplasty or submucous resection, with or without cartilage scoring, contouring or replacement with graft
30620	Septal or other intranasal dermatoplasty (does not include obtaining graft)

Applicable Diagnosis Codes

C30.0	Malignant neoplasm of nasal cavity
-------	------------------------------------

C31.0	Malignant neoplasm of maxillary sinus
C31.1	Malignant neoplasm of ethmoidal sinus
C31.2	Malignant neoplasm of frontal sinus
C31.3	Malignant neoplasm of sphenoid sinus
C31.8	Malignant neoplasm of overlapping sites of accessory sinuses
C41.0	Malignant neoplasm of bones of skull and face
C79.89	Secondary malignant neoplasm of other specified sites
C79.9	Secondary malignant neoplasm of unspecified site
D14.0	Benign neoplasm of middle ear, nasal cavity and accessory sinuses
D16.4	Benign neoplasm of bones of skull and face
J32.0	Chronic maxillary sinusitis
J32.1	Chronic frontal sinusitis
J32.2	Chronic ethmoidal sinusitis
J32.3	Chronic sphenoidal sinusitis
J32.4	Chronic pansinusitis
J32.8	Other chronic sinusitis
J32.9	Chronic sinusitis, unspecified
J33.0	Polyp of nasal cavity
J33.1	Polypoid sinus degeneration
J33.8	Other polyp of sinus
J33.9	Nasal polyp, unspecified
J34.0	Abscess, furuncle and carbuncle of nose
J34.1	CYST and mucocele of nose and nasal sinus
J34.2	Deviated nasal septum
M95.0	Acquired deformity of nose
Q30.3	Congenital perforated nasal septum
R04.0	Epistaxis
R09.81	Nasal congestion

References

Dinis PB, Haider H. Septoplasty: long-term evaluation of results. *Am J Otolaryngol.* 2002;23:85-90.

Doyle RL. *Healthcare Management Guidelines. Vol 1. 2nd ed.* Seattle: Milliman and Robertson; 1997.

Godley FA, Nemeroff RF, Josephson JS. Current trends in rhinoplasty and the nasal airway. *Med Clin North Am.* 1993;77:643-656.

Grossman ZD, Katz DS, Santelli ED, et al. Sinusitis. In: *Cost-Effective Diagnostic Imaging: The Clinician's Guide. 3rd ed.* St. Louis: Mosby; 1995:376-377.

Huerter JV. Functional endoscopic sinus surgery and allergy. *Otolaryngol Clin North Am.* 1992;25:231-238.

Septoplasty

Last review: August 11, 2017

Page 4 of 4

Lund VJ. Office evaluation of nasal obstruction. *Otolaryngol Clin North Am.* 1992;25:803-815.

Marshall, KG, Elhamy LA. *Disorders of the Nose and Paranasal Sinuses.* Littleton, Mass: PSG Publishing Co.; 1987.

Schuller DE, Schleuning AJ. DeWeese and Saunders' *Otolaryngology: Head and Neck Surgery.* 8th ed. St. Louis: Mosby; 1994.

Specialty-matched clinical peer review.