



# Reimbursement Policy:

## Evaluation of Dry Eyes - Lab Benefit Program (LBM)

POLICY NUMBER	EFFECTIVE DATE:	APPROVED BY
AHS-G2138	3/01/2023	RPC (Reimbursement Policy Committee)

**Reimbursement Guideline Disclaimer:** We have policies in place that reflect billing or claims payment processes unique to our health plans. Current billing and claims payment policies apply to all our products, unless otherwise noted. We will inform you of new policies or changes in policies through postings to the Reimbursement Policies webpage on emblemhealth.com. Further, we may announce additions and changes in our provider manual and/or provider newsletters which are available online and emailed to those with a current and accurate email address on file. The information presented in this policy is accurate and current as of the date of this publication.

The information provided in our policies is intended to serve only as a general reference resource for services described and is not intended to address every aspect of a reimbursement situation. Other factors affecting reimbursement may supplement, modify or, in some cases, supersede this policy. These factors may include, but are not limited to, legislative mandates, physician or other provider contracts, the member's benefit coverage documents and/or other reimbursement, and medical or drug policies. Finally, this policy may not be implemented the same way on the different electronic claims processing systems in use due to programming or other constraints; however, we strive to minimize these variations.

We follow coding edits that are based on industry sources, including, but not limited to, CPT® guidelines from the American Medical Association, specialty organizations, and CMS including NCCI and MUE. In coding scenarios where there appears to be conflicts between sources, we will apply the edits we determine are appropriate. We use industry-standard claims editing software products when making decisions about appropriate claim editing practices. Upon request, we will provide an explanation of how we handle specific coding issues. If appropriate coding/billing guidelines or current reimbursement policies are not followed, we may deny the claim and/or recoup claim payment.

[POLICY DESCRIPTION](#) | [INDICATIONS AND/OR LIMITATIONS OF COVERAGE](#) | [DEFINITIONS](#) | [SCIENTIFIC BACKGROUND](#) | [GUIDELINES AND RECOMMENDATIONS](#) | [APPLICABLE STATE AND FEDERAL REGULATIONS](#) | [APPLICABLE CPT/HCPCS PROCEDURE CODES](#) | [EVIDENCE-BASED SCIENTIFIC REFERENCES](#) | [REVISION HISTORY](#)

### Policy Description:

Dry eye disease (dysfunctional tear syndrome, DED) is defined by the Dry Eye Workshop II as “a multifactorial disease of the ocular surface characterized by a loss of homeostasis of the tear film, and accompanied by ocular symptoms, in which tear film instability and hyperosmolarity, ocular surface inflammation and damage, and neurosensory abnormalities play etiological roles.”<sup>1</sup> Five to fifteen percent of the United States population suffers from dry eye disease, leaving a substantial burden on functional vision, general health status, and workplace productivity.<sup>2</sup>

### Indications and/or Limitations of Coverage:

Application of coverage criteria is dependent upon an individual's benefit coverage at the time of the request. Specifications pertaining to Medicare and Medicaid can be found in the Applicable State and Federal Regulations section of this policy document.

- 1) For individuals suspected of having dry eye, testing of tear osmolarity **MEETS COVERAGE CRITERIA** in **any** of the following situations:
  - a) To help determine the severity of dry eye disease.
  - b) To monitor effectiveness of therapy.
- 2) For individuals suspected of having dry eye disease based on comprehensive eye examination, testing for MMP-9 protein in human tears **DOES NOT MEET COVERAGE CRITERIA**.

## Reimbursement Policy:

### Evaluation of Dry Eyes - Lab Benefit Program (LBM)

- 3) For individuals suspected of having dry eye disease, testing for lactoferrin and/or IgE **DOES NOT MEET COVERAGE CRITERIA.**

*The following does not meet coverage criteria due to a lack of available published scientific literature confirming that the test(s) is/are required and beneficial for the diagnosis and treatment of an individual's illness.*

- 4) For individuals suspected of having dry eye disease, all other testing not discussed above **DOES NOT MEET COVERAGE CRITERIA.**

### Definitions:

Term	Definition
AAO	American Academy of Ophthalmology
AAOPT	American Academy of Optometry
AOA	American Optometric Association
ASCRS	American Society of Cataract and Refractive Surgery
ATD	Advanced tear diagnostics
CA-6	Carbonic anhydrase-6
CLIA '88	Clinical Laboratory Improvement Amendments of 1988
CMS	Centers for Medicare and Medicaid Services
DED	Dry eye disease
DEWS	Dry eye workshop
DTS	Dysfunctional tear syndrome
FDA	Food and Drug Administration
IgA	Immunoglobulin A
IgE	Immunoglobulin E
IgG	Immunoglobulin G
IgM	Immunoglobulin M
LASIK	Laser-assisted in situ keratomileusis
LDTs	Laboratory-developed tests
MMP	Matrix metalloproteinase
MMP-9	Matrix metalloproteinase-9
NIKBT	Non-invasive tear breakup time
OSD	Ocular surface disorders
OSDI	Ocular surface disease index

## Reimbursement Policy:

### Evaluation of Dry Eyes - Lab Benefit Program (LBM)

Term	Definition
OSS	Ocular surface staining
PSP	Parotid secretory protein
SP-1	Salivary protein-1
SPEED	Standard patient evaluation of eye dryness
TBUT	Tear break-up time
TFBUT	Tear film break-up time
TFOS	Tear film & ocular surface
VAS	Visual analogue scale

### Scientific Background:

Tears are necessary for maintaining the health of the inner and outer surfaces of the eyelid and for providing clear vision. The tear film of the eye consists of aqueous, mucous, and lipid components. A healthy tear film is necessary for protecting and moisturizing the cornea, as well as for providing a refracting surface for light entering the eye.<sup>3</sup> Dysfunction of any component of the tear film can lead to dry eye disease (dysfunctional tear syndrome, DED). Dry eye is a common and often chronic problem, particularly in older adults as age affects the entire lacrimal functional unit.<sup>4</sup> The exact prevalence of dry eye is unknown due to difficulty in defining the disease and the lack of a single diagnostic test to confirm its presence, but the 2013 National Health and Wellness Survey estimated the rate of dry eye in the United States to be 6.8%, or about 16.4 million people; prevalence tended to increase with age, with the 18-34 age group only comprising 2.7% of the total and the 75+ age group comprising 18.6%.<sup>5,6</sup> Risk factors for dry eye include increasing age, systemic comorbidities such as diabetes and autoimmune disease, and therapeutic treatments for anxiety, depression, and sleep disorders.<sup>7</sup>

Further, the 2017 Tear Film & Ocular Surface (TFOS) Society International Dry Eye Workshop (DEWS) II reported that “the core mechanism of dry eye disease is tear hyperosmolarity, which is the hallmark of the disease.”<sup>1</sup>

Dry eye is classified into two general groups: decreased tear production and increased evaporative loss. Decreased tear production may lead to hyperosmolarity of the tear film and inflamed ocular surface cells. An age-related ductal obstruction is the most common cause of decreased tear production. Increased evaporative loss is typically caused by problems in the meibomian gland when the glands that produce the lipid portion of the tear film fail. This lipid portion normally allows the tear film to spread evenly, minimizing evaporation. In both groups, tear film hyperosmolarity and subsequent ocular surface inflammation lead to the variety of symptoms and signs associated with dry eye.<sup>6</sup>

Most patients will present with symptoms of chronic eye irritation, such as red eyes, light sensitivity, blurred vision, or unusual sensations (gritty, burning, foreign, etc.). However, significant variability in the patient-reported symptoms and signs, as well as a lack of correlation between these symptoms and signs, make it difficult to diagnose dry eye, and no single definitive test to diagnose dry eye exists. Dry eye is typically diagnosed with a combination of patient symptoms and physical findings, such as reduced blink rate or eyelid malposition.<sup>6</sup> Incomplete blinking may also be considered for mild-to-moderate dry eye assessment.<sup>8</sup> Further, visual acuity was found to be particularly poor in those with vision-related symptoms due to dry eyes.<sup>9</sup>

## Reimbursement Policy:

### Evaluation of Dry Eyes - Lab Benefit Program (LBM)

The primary way to treat dry eye is artificial tears, although corticosteroids, topical cyclosporine A, or anti-inflammatories such as Lifitegrast ophthalmic solution five percent may be used to supplement treatment. Avoiding environmental factors, such as heavy smoke or dry heating air, is also recommended.<sup>10</sup> It was recently reported by Holland, et al. (2019), who reviewed two decades worth of data on the safety and efficacy of controlled topical ophthalmic drug administration for DED treatment, that poor standardization of endpoints across studies causes challenges in the improvement of this field. However, recent advances in drug delivery and a greater understanding of DED will assist in the improvement of ophthalmic drugs.

Accurate diagnosis of dry eye disease requires a variety of tests including patient-reported symptom questionnaires, tear film break-up time (TFBUT), Schirmer test, ocular surface staining, and meibomian gland functionality. However, many of these tests lack consistency and reliability in diagnosis. New tools have been developed which allow for the quantification of tear film characteristics including measurement of tear osmolarity and measurement of inflammatory mediators such as matrix metalloproteinase enzymes, and biomarkers such as lactoferrin.<sup>6</sup>

#### *Tear Osmolarity*

Osmolarity is a measurement of the concentration of dissolved solutes in a solution. Hyperosmolarity of the tear film is a recognized and validated marker of dry eye. The following tear osmolarity thresholds have been suggested for establishing the severity of dry eyes: 270-308 mOsm/L for normal eyes, 308-316 mOsm/L for mild dry eye, and >316 mOsm/L for moderate to severe dry eye.<sup>12</sup> Tomlinson, et al. (2006) suggested a cut-off of 316 mOsm/L, but the sensitivity was found to be 0.59 when applied to the independent sample described in the study. Furthermore, decreasing the cut-off to increase the sensitivity decreased the specificity and overall accuracy significantly. Overall, the overlap between normal and dry eyes contributes heavily to the difficulty in establishing a cut-off.<sup>13</sup> Some studies suggest that osmolarity shows the strongest correlation with severity of dry eye based on the metrics used, but at the same time lack correlation to other objective signs of dry eye. In general, tear osmolarity results vary between clinical signs and symptoms, which can make them difficult to interpret.<sup>14</sup>

The test “TearLab” is based on assessment of the osmolarity of tears. TearLab collects a 50 µL tear sample, analyzes its electrical impedance, and provides an assessment of the osmolarity of the sample and thereby the tear.<sup>3</sup> Baenninger, et al. (2018) completed an extensive systematic review investigating 1362 healthy eyes of participants from 33 different studies; this review found a weighted mean osmolarity of 298 mOsm/L via the TearLab test. Final comments from the researchers highlighted the great variability of osmolarity measurements that were found with the TearLab system, suggesting caution when interpreting TearLab osmolarity results.<sup>15</sup>

#### *Matrix Metalloproteinase (MMP) Enzymes*

Inflammation is a common factor across the subtypes of DED. Levels of inflammatory mediators, such as cytokines, may be assessed in the tear film. For example, the matrix metalloproteinase (MMP) enzymes play an important role in wound healing and inflammation by degrading collagen. Elevated levels of MMP-9, a member of the MMP family produced by corneal epithelial cells<sup>16,17</sup>, have been observed in the tears of patients with dry eye.<sup>18</sup> A study with 101 patients with DED and controls (54 controls, 47 with DED) was performed to assess correlation of the protein MMP-9 with dry eye. All 101 underwent MMP-9 testing of the tear film and were evaluated for symptoms and signs of DED. The tear film was then analyzed for MMP-9 by InflammDry, which detects MMP-9 levels of more than 40 ng/mL. The MMP-9 results were positive in 19 of the 47 dry eye patients (40.4%) and three of the 54 controls (5.6%). The authors concluded that “MMP-9 correlated well with other dry eye tests and identified the presence of ocular surface inflammation in 40% of confirmed dry eye patients,” and suggested it may be helpful to identify patients with autoimmune disease and ocular surface inflammation.<sup>19</sup> The American Academy of Ophthalmology (AAO) has noted MMP-9 does not differentiate dry eye from any other inflammatory ocular surface disease and does not include this test in its appendix on diagnostic tests.<sup>14</sup>

## Reimbursement Policy:

### Evaluation of Dry Eyes - Lab Benefit Program (LBM)

#### *Lactoferrin*

Another biomarker associated with inflammation is lactoferrin. Lactoferrin is thought to promote the healing process resulting from inflamed dry eyes and is used to assess the lacrimal glands.<sup>3</sup> “Lactoferrin & Dry Eye Disease (DED): The measurement of ocular lactoferrin, as a biomarker, has long been established and accepted as the “medical standard” in assessing the secretory function of the lacrimal gland. Low lactoferrin levels directly correlate to aqueous deficiency. Normal lactoferrin levels indicate normal lacrimal gland function.”<sup>20</sup> The AXIM Eye Lactoferrin test has a specificity of 98%, and a sensitivity of 83%. A High Lf: > 1.0 mg/ml and a Low Lf: < 0.8 mg/ml.<sup>20</sup>

#### *Additional Tests*

Other tests noted by the American Academy of Optometry (AAOPT) are the tear break-up time test, the ocular surface dry staining test, the Schirmer test, and the fluorescein dye disappearance test. The tear break-up time test evaluates the precorneal tear film’s stability with a fluorescein dye, which is inserted in the lower eyelid. If the tear film layer develops a dark discontinuity (usually blue) in under ten seconds, the result is considered abnormal. The ocular surface dry staining test stains areas of discontinuity of the corneal epithelial surface, which may contribute to dryness. A fluorescein dye is typically used, although a rose bengal dye or a lissamine green dye may be used as well. The Schirmer test quantifies the amount of tears produced by each eye. This is done by placing small strips of filter paper in the lower eyelid and checking the length (in mm) of wet strips in a certain amount of time. This test is noted as an extremely variable test, so it should not be used as the only diagnostic test. Finally, the fluorescein dye disappearance test places a certain amount of fluorescein dye on the ocular surface, and then evaluates how much of that dye was cleared from the surface.<sup>6,14</sup>

Evaluation of dry eyes is difficult for numerous reasons. Currently, no “gold standard” or globally accepted guideline for diagnosis of dry eye exists, and no threshold between healthy and affected eyes has been established. Many other features of testing (repeatability, high variability, including highly variable sensitivity and specificity of tests and dependence on clinical conditions) and the disease itself—its multifactorial status, examiner subjectivity, reliance on patient-based questionnaires, for example—make diagnosis of dry eye especially challenging.<sup>21</sup> Despite promising sensitivities, specificities, or other strong statistical findings, these numbers should still be considered in the context of clinical findings.<sup>14</sup>

#### ***Clinical Utility and Validity***

##### *Tear Osmolarity*

Brissette, et al. (2019) measured the utility of the TearLab test in 100 patients with DED-like symptoms who had normal tear osmolarity results. This study aimed to use the test to identify diagnoses other than DED. All patients included in the study had a normal tear osmolarity test (<308 mOsm/L in each eye, and an inter-eye difference < eight mOsm/L). The researchers report that “A possible alternate diagnosis was established in 89% of patients with normal tear osmolarity testing. The most frequent diagnoses included anterior blepharitis (26%) and allergic conjunctivitis (21%).”<sup>22</sup> This highlights the utility of the TearLab test to differentiate between DED and other eye disorders with overlapping symptoms.

In a retrospective study by Tashbayev, et al. (2020), 757 patients diagnosed with symptomatic DED were recruited to investigate the clinical utility of tear osmolarity measurement. The TearLab osmometer was used to measure osmolarity in both eyes and the results were compared to Ocular Surface Disease Index (OSDI), TFBUT, ocular surface staining (OSS), Schirmer test, and meibomian gland functionality tests. According to their data, TearLab results were not significantly different between the healthy controls and the DED patients. Many studies confirm that tear osmolarity greater than 308 MOsm/mL indicates a loss of homeostasis in the tear, therefore, is used as a cut-off value. Many of the healthy controls had tear osmolarity levels above the

## Reimbursement Policy:

### Evaluation of Dry Eyes - Lab Benefit Program (LBM)

recommended cut-off value of 308 mOsm/L, and a substantial proportion of the diagnosed DED patients had tear osmolarity levels below the cut-off value. In the DED patient group, osmolarity levels in the right and left eye were 275–398 mOsm/L and 272–346 mOsm/L, respectively. In the control group, osmolarity levels in the right and left eyes were 281–369 mOsm/L and 275–398 mOsm/L, respectively. Therefore, the authors suggest that "tear osmolarity measured with TearLab osmometer cannot be used as a key indicator of DED."<sup>23</sup>

As shown in the above studies, there have been issues in the past regarding the use of tear osmolarity as a diagnostic tool. First, no criteria for the measurement of osmolarity have been established. Studies reviewing osmolarity as a diagnostic tool do not use uniform numbers in their calculations (i.e., no uniform cut-off values, no standardized severity measures, etc.). To compound this issue, high variance in osmolarity due to outside factors, such as sleep deprivation, altitude, or even whether the right or left eye was used to produce the tears, can occur. This difficulty in establishing osmolarity ranges has caused an overlap between the ranges of healthy and dry eye osmolarity. Although measuring fluctuations between osmolarity readings has been suggested as a diagnostic (caused by increased instability), the line between healthy eyes and dry eyes is blurred.<sup>3</sup> However, a recent report by the TFOS DEWS II states that tear osmolarity "is a global, early stage marker of the disease and has been shown to be able to effectively track therapeutic response and inform the clinician as to whether there has been a loss of tear film homeostasis."<sup>1</sup>

#### *MMP Enzymes*

Chan, et al. (2016) aimed to assess the utility of MMP-9 measurement in patients with post-laser-assisted in situ keratomileusis (LASIK) dry eyes compared to aged-matched controls. The *InflammaDry* was used to measure MMP-9 levels in tear film. Results showed that "The tear film MMP-9 levels were  $52.7 \pm 32.5$  ng/mL in dry eyes and  $4.1 \pm 2.1$  ng/mL in normal eyes ( $p < 0.001$ ). MMP-9 levels were  $>40$  ng/mL in seven out of 14 (50.0%) post-LASIK dry eyes. The *InflammaDry* was positive in eight out of 14 (57.1%) post-LASIK eyes. All positive cases had tear film MMP-9 levels  $\geq 38.03$  ng/mL. Agreement between *InflammaDry* and MMP-9 was excellent with Cohen  $\kappa$  value of 0.857 in post-LASIK dry eyes."<sup>24</sup> However, only half of the post-LASIK patients with dry eyes exhibited significant inflammation with heightened levels of MMP-9.<sup>24</sup>

A cross-sectional study by Jun JH (2020) investigated if the tear volume in dry eye disease (DED) patients affects the results of the MMP-9 immunoassay (*InflammaDry*). A total of 188 DED patients were enrolled in the study. Positive MMP-9 tests were confirmed in 120 patients, and negative results were noted in 68 patients. However, the authors observed that with a small sample volume, the reliability of the test result was impaired. The manufacturer also pointed out that less than six  $\mu$ l of sample volume could produce false-negative results. In this study, patients with higher tear volumes showed higher band densities, but subjects with lower tear volumes showed lower band densities on the immunoassay. In conditions such as Sjögren syndrome that present with markedly decreased tear secretion, *InflammaDry* could display negative results despite the elevated tear MMP-9 concentration. In addition, "among the participants of the present study, a strong positive band was identified even in patients with mild or nearly no fluorescein staining of the cornea and conjunctiva, who are expected to have very mild inflammatory eye surface inflammation."<sup>25</sup> In conclusion, this study determined the volume dependency of the MMP-9 immunoassay, which could induce false-negative results clinically.<sup>25</sup>

Lee, et al. (2021) conducted a cross-sectional study to analyze the association of MMP-9 immunoassay results with the severity of DED symptoms and signs. Using 320 patients, the researchers evaluated the clinical signs based on the OSDI score, visual analogue scale (VAS), TBUT, "tear volume evaluation by tear meniscometry, and staining scores of the cornea and conjunctiva by the Oxford grading scheme." They found that "positive MMP-9 immunoassay results were significantly related to shorter tBUT,  $tBUT \leq$  three seconds, higher corneal staining score, corneal staining score  $\geq$  two, and conjunctival staining score  $\geq$  two" which indicated a worsening severity of ocular signs in DED. The researchers also performed semiquantitative analyses, basing the reagent band density on a four-point scale ranging from negative (zero) to strongly positive (three), and found that these

## Reimbursement Policy:

### Evaluation of Dry Eyes - Lab Benefit Program (LBM)

results positively correlated with higher corneal staining scores and negatively correlated with TBUT. However, despite these correlating results, the researchers found that their quantitative analysis, which would've been the most accurate way to evaluate tear MMP-9 levels, yielded no correlation between "immunoassay band density and the clinical signs and symptoms of DE." This likely indicates the need for more studies with less selection bias and greater consideration of DED subtypes, as this finding was contrary to established literature.

Choi, et al. (2023) conducted a study comparing positive MMP-9 presence against an increased tear osmolarity measurement to diagnose severity of DED. The researchers found that those who tested positive for MMP-9 via immunoassay had "higher corneal fluorescein staining score and worse DED severity," as well as a worse ocular surface staining score with statistical significance. In using a cutoff for tear osmolarity level of 308 mOSm/L, they found "no significant difference in dry eye signs and symptoms," but "higher tear osmolarity was associated with ocular surface staining score in patients with severe DED and [evaporative dry eye]." Though historically tear osmolarity has been found to be useful in diagnosing DED, the researchers attribute the difference in their current findings to the study population, and that it may be better for identifying clinical severity in those already diagnosed with DED by other means.<sup>27</sup>

#### *Lactoferrin*

A meta-analysis was performed to highlight the potential role of tear lactoferrin as a diagnostic biomarker for DED. All original studies reporting an estimate of the average lactoferrin concentration in healthy subjects and those affected by DED were searched. A pooled mean difference of 0.62 (95% CI, 0.35–0.89) in lactoferrin concentration was observed in DED patients, showing a significant decrease in lactoferrin concentrations in the tears of subjects affected by DED. A study reported that administration of lactoferrin protein in mice led to a decrease in oxidative damage and an enhancement of tear function.<sup>28</sup> Lastly, the author notes that "to compare data across studies and to validate lactoferrin as a diagnostic biomarker, there is still a need for further development of standardized protocols of tear collection, processing and storage."<sup>29</sup>

## Guidelines and Recommendations:

### Dysfunctional Tear Syndrome (DTS) Panel

A study assessed the new diagnostic techniques and treatment options for DED and associated tear film disorders. Experts from the Cornea, External Disease, and Refractive Society (DTS Panel) convened by the study found examining tear osmolarity useful in diagnosis "in combination with other clinical assessments and procedures." The same panel also stated that the use of MMP-9 may only be valid for more severe cases of dry eye since the diagnostic test is only positive past 40 ng/mL. The panel recommended that osmolarity be evaluated before any ocular surface assessment, then an evaluation of ocular inflammation can be done, and finally a Schirmer strip test should be done.<sup>12</sup>

### American Academy of Ophthalmology (AAO)

The AAO states "no single test is adequate for establishing the diagnosis of dry eye" and recommends that the combination of findings from diagnostic tests can be useful to understanding a patient's condition. In particular, the AAO states, "tests results should be considered within the context of symptoms and other clinical findings." This statement was reaffirmed in the dry eye syndrome preferred practice pattern guidelines of 2023.<sup>30</sup> "Pharmacological and procedural treatments are associated with improvements in patient symptoms and clinical signs, although chronic therapy and patient compliance are necessary in most instances. Topical cyclosporine treatment has long been used in the treatment of dry eye and shown to have clinical benefits. Topical

## Reimbursement Policy:

### Evaluation of Dry Eyes - Lab Benefit Program (LBM)

cyclosporine, in some instances, leads to long-term treatment-free remission of patient symptoms and signs. Lifitegrast is a lymphocyte function-associated antigen-1 antagonist developed to treat dry eye syndrome (also known as dry eye disease), but the exact mechanism of action of lifitegrast in dry eye is unknown. Topical lifitegrast five percent has been approved by the US Food and Drug Administration for treatment of dry eye.”<sup>14</sup> In their 2022 summary benchmarks for the cornea/external disease preferred practice pattern guidelines, the AAO cites that diagnostic tests may include “tear break-up time, ocular surface dye staining, Schirmer test, fluorescein dye disappearance test/tear function index, and tear osmolarity test.”<sup>31</sup>

Rather than relying solely on a single measure of tear osmolarity, correlation with clinical findings or differences in osmolarity over time or under different conditions is more informative for confirming the diagnosis of dry eye. Indeed, most recent studies confirm that normal subjects have exceptionally stable tear film osmolarity, whereas tear osmolarity values in dry eye subjects become unstable quickly and lose homeostasis with environmental changes. These data reinforce the long-held belief that tear film instability due to increased evaporation of tears resulting in hyperosmolarity (i.e., evaporative dry eye) is a core mechanism of the disease.”<sup>14</sup> The guideline covers the currently used diagnostic tests, which are as follows: assessment of tear osmolarity, MMP-9, tear production, fluorescein dye or tear function index, tear break up time, ocular surface dye staining, and lacrimal gland function.<sup>14</sup> The following table is provided by Akpek, et al. (2019):

The following table is provided by Akpek et al. (2019):

**Table 2:** Characteristic Findings for Dry Eye Disease Diagnostic Tests

Test	Characteristic Findings
<b>Tear osmolarity</b>	Elevated; test-to-test variability; inter-eye differences considered abnormal
<b>Matrix metalloproteinase-9</b>	Indicates presence of inflammation which dictates treatment
<b>Aqueous tear production (Schirmer test)</b>	10 mm or less considered abnormal
<b>Fluorescein dye disappearance test/tear function test</b>	Test result is compared with a standard color scale
<b>Tear break-up time</b>	Less than 10 seconds considered abnormal
<b>Ocular surface dye staining</b>	Staining of inferior cornea and bulbar conjunctiva typical
<b>Lacrimal gland function</b>	Decreased tear lactoferrin concentrations

### Tear Film & Ocular Surface (TFOS) Society

The TFOS society held the International Dry Eye Workshop II in 2017. From this workshop, the society published recommendations on the management and treatment of DED. The authors state that when diagnosing DED, it is important to distinguish between the type (aqueous deficient dry eye or evaporative dry eye) and to determine the underlying etiology as this is crucial for proper management.<sup>32</sup> These guidelines also stated that “neurotrophic keratopathy accompanied by neuropathic pain and symptoms should definitely be considered in differential diagnosis of patients with intense symptoms despite mild signs.”<sup>32</sup>

Regarding diagnostic testing, the TFOS states that any patient who obtains a positive score on the Dry Eye Questionnaire-5 or Ocular Surface Disease Index should be subject to a clinical examination. “The presence of any one of three specified signs; reduced non-invasive break-up time; elevated or a large interocular disparity in osmolarity; or ocular surface staining (of the cornea, conjunctiva or lid margin) in either eye, is considered representative of disrupted homeostasis, confirming the diagnosis of DED. If a patient has DED symptoms and

## Reimbursement Policy:

### Evaluation of Dry Eyes - Lab Benefit Program (LBM)

their practitioner does not have access to all these tests, a diagnosis is still possible, based on a positive result for any one of the markers, but may require referral for confirmation if the available homeostasis markers are negative.”<sup>32</sup> After confirmation with any of the aforementioned tests (i.e. reduced non-invasive break-up time less than ten seconds, an elevated or large interocular disparity in osmolarity  $\geq 308$  mOsm/L in either eye or an interocular difference greater than eight mOsm/L, or ocular surface staining including greater than five corneal spots, greater than nine conjunctival spots, or a lid margin  $\geq$  two mm in length and  $\geq 25\%$  in width), further evaluation should be conducted including meibography, lipid interferometry, and tear volume measurement to assess severity and help determine the best treatment plan.<sup>32</sup>

Further, the consensus recommendation from the society on tear osmolarity testing states, “The low variation of normal subjects contributes to the high specificity of the marker and makes it a good candidate for parallelization and therapeutic monitoring. Accordingly, normal subjects don't display elevated osmolarity, so a value over 308 mOsm/L in either eye or a difference between eyes greater than eight mOsm/L are good indicators of a departure from tear film homeostasis and represent a diseased ocular surface.”<sup>1</sup>

Regarding MMP-9 testing, the guidelines state that “With the availability of newer immunosuppressive medications and trials concerning these drugs it is logical that inflammation should be assessed. The exact modality used may need to be varied depending on the pathway or target cell upon which the immunosuppressive drug acts, and such diagnostic tools should be used for refining patient selection as well as monitoring after commencement of treatment. Costs of these diagnostic tests should be considered, but these should be calculated from a holistic standpoint. For example, if the tests can assist the channeling of patients to appropriate healthcare services there may be cost savings for reduced referrals.”<sup>1</sup>

#### American Optometric Association

The AOA published consensus-based clinical practice guidelines for care of a patient with ocular surface disorders. These guidelines note that there is a “lack of a defined diagnostic test or protocol and a lack of congruity between patient symptoms and clinical tests.” The AOA also notes that the condition itself is ill defined and that dry eye is often a symptom of another condition such as blepharitis or another glandular dysfunction.<sup>33</sup> There have not been any updates on this topic from the AOA since this 2010 statement.

#### Consensus Guidelines for Management of Dry Eye Associated with Sjögren Disease

In 2015, clinical guidelines for management of dry eye associated with Sjögren disease were published by a consensus panel which evaluated reported treatments for DED. The recommendations state, “Evaluation should include symptoms of both discomfort and visual disturbance as well as determination of the relative contribution of aqueous production deficiency and evaporative loss of tear volume. Objective parameters of tear film stability, tear osmolarity, degree of lid margin disease, and ocular surface damage should be used to stage severity of dry eye disease to assist in selecting appropriate treatment options. Patient education with regard to the nature of the problem, aggravating factors, and goals of treatment is critical to successful management. Tear supplementation and stabilization, control of inflammation of the lacrimal glands and ocular surface, and possible stimulation of tear production are treatment options that are used according to the character and severity of dry eye disease.”<sup>34</sup> Further, tear osmolarity was identified as the testing method with the highest level of evidence for all DED related tests.

#### American Society of Cataract and Refractive Surgery (ASCRS) Cornea Clinical Committee

American Society of Cataract and Refractive Surgery (ASCRS) released guidelines to aid surgeons in diagnosing visually significant ocular surface disorders (OSD) before refractive surgery. The ASCRS Cornea Clinical Committee recommends initial screening procedures including ASCRS Standard Patient Evaluation of Eye Dryness (SPEED) II questionnaire, tear osmolarity, and matrix metalloproteinase (MMP-9) testing. If any

## Reimbursement Policy:

### Evaluation of Dry Eyes - Lab Benefit Program (LBM)

of the three initial screening tests are abnormal, the patient is at risk for ocular surface disease, and additional diagnostic tests can be performed to determine dry eye sub-type. Non-invasive tests are recommended to minimize disruption to the ocular surface, cornea, and tear film. These tests include tear lipid layer thickness, noninvasive tear breakup time (NIKBUT), tear meniscus height, meibography, topography, tear lactoferrin levels, and measures of optical scatter. However, these tests are not essential to the fundamental algorithm.

The ASCRS also notes a point of care test that assesses lactoferrin levels (TearScan). The guideline notes its three proprietary biomarkers which are as follows: “salivary protein-1 (SP-1, immunoglobulin A [IgA], immunoglobulin G [IgG], immunoglobulin M [IgM]); (2) carbonic anhydrase-6 (CA-6, IgA, IgG, IgM); and (3) parotid secretory protein (PSP, IgA, IgG, IgM)”. The authors comment that this test can be used to detect Sjögren syndrome early. However, the authors also note that “no member of the ASCRS Cornea Clinical Committee has used it [TearScan] in clinical practice.”<sup>35</sup>

### Applicable State and Federal Regulations:

DISCLAIMER: If there is a conflict between this policy and any relevant, applicable government policy for a particular member [e.g., Local Coverage Determinations (LCDs) or National Coverage Determinations (NCDs) for Medicare and/or state coverage for Medicaid], then the government policy will be used to make the determination. For the most up-to-date Medicare policies and coverage, please visit the Medicare search website: <https://www.cms.gov/medicare-coverage-database/search.aspx>. For the most up-to-date Medicaid policies and coverage, visit the applicable state Medicaid website.

### Food and Drug Administration (FDA)

On December 3, 1993, the FDA approved the lactoferrin microassay system by Touch Scientific, Inc.<sup>36</sup> Lactoferrin diagnostic kits are commercially available options for tear film biomarkers.<sup>3</sup>

On May 14, 2009, the FDA approved the TearLab Osmolarity System created by Ocusense Inc. From the FDA site: this device is used “to measure the osmolality of human tears to aid in the diagnosis of patients with signs or symptoms of DED, in conjunction with other methods of clinical evaluation.”<sup>37</sup> In 2022, Ocusense Inc. rebranded as Trukera Medical and continues to manage the TearLab Osmolarity System.<sup>38</sup>

On November 20, 2013, the FDA approved InflammDry created by Rapid Pathogen Screening Inc. From the FDA site: “InflammDry is a rapid, immunoassay test for the visual, qualitative in vitro detection of elevated levels of the MMP-9 protein in human tears from patients suspected of having dry eye to aid in the diagnosis of dry eye in conjunction with other methods of clinical evaluation. This test is intended for prescription use at point-of-care sites.”<sup>39</sup>

Many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). As an LDT, the U. S. Food and Drug Administration has not approved or cleared this test; however, FDA clearance or approval is not currently required for clinical use.

## Reimbursement Policy:

### Evaluation of Dry Eyes - Lab Benefit Program (LBM)

#### Applicable CPT/HCPCS Procedure Codes:

CPT	Code Description
82785	Gammaglobulin (immunoglobulin); IgE
83516	Immunoassay for analyte other than infectious agent antibody or infectious agent antigen; qualitative or semiquantitative, multiple step method
83520	Immunoassay for analyte other than infectious agent antibody or infectious agent antigen; quantitative, not otherwise specified
83861	Microfluidic analysis utilizing an integrated collection and analysis device, tear osmolarity

Current Procedural Terminology© American Medical Association. All Rights reserved.

Procedure codes appearing in Medical Policy documents are included only as a general reference tool for each policy. They may not be all-inclusive.

#### Evidence-based Scientific References:

1. Craig JP, Nichols KK, Alpek MD, et al. TFOS DEWS II Definition and Classification Report. *The ocular surface*. Oct 2017;15(4):276-283. doi:10.1016/j.jtos.2017.08.003
2. Dana R, Meunier J, Markowitz JT, Joseph C, Siffel C. Patient-Reported Burden of Dry Eye Disease in the United States: Results of an Online Cross-Sectional Survey. *Am J Ophthalmol*. Aug 2020;216:7-17. doi:10.1016/j.ajo.2020.03.044
3. Willcox MDP, Argüeso P, Georgiev GA, et al. TFOS DEWS II Tear Film Report. *The ocular surface*. Jul 2017;15(3):366-403. doi:10.1016/j.jtos.2017.03.006
4. Ezuddin NS, Alawa KA, Galor A. Therapeutic Strategies to Treat Dry Eye in an Aging Population. *Drugs & aging*. Jul 2015;32(7):505-13. doi:10.1007/s40266-015-0277-6
5. Farrand KF, Fridman M, Stillman IO, Schaumberg DA. Prevalence of Diagnosed Dry Eye Disease in the United States Among Adults Aged 18 Years and Older. *Am J Ophthalmol*. Oct 2017;182:90-98. doi:10.1016/j.ajo.2017.06.033
6. Shtein R. Dry eye disease. Wolters Kluwer. 11/16, 2018. Updated Aug 07, 2024. <https://www.uptodate.com/contents/dry-eye-disease>
7. Periman. The Immunological Basis of Dry Eye Disease and Current Topical Treatment Options. *Journal of Ocular Pharmacology and Therapeutics*. 2020;36(3):137-146. doi:10.1089/jop.2019.0060
8. Jie Y, Sella R, Feng J, Gomez ML, Afshari NA. Evaluation of incomplete blinking as a measurement of dry eye disease. *The ocular surface*. Jul 2019;17(3):440-446. doi:10.1016/j.jtos.2019.05.007
9. Szczotka-Flynn LB, Maguire MG, Ying GS, et al. Impact of Dry Eye on Visual Acuity and Contrast Sensitivity: Dry Eye Assessment and Management Study. *Optom Vis Sci*. Jun 2019;96(6):387-396. doi:10.1097/OPX.0000000000001387
10. Messmer EM. The pathophysiology, diagnosis, and treatment of dry eye disease. *Deutsches Arzteblatt international*. Jan 2015;112(5):71-81; quiz 82. doi:10.3238/arztebl.2015.0071
11. Holland EJ, Darvish M, Nichols KK, Jones L, Karpecki PM. Efficacy of topical ophthalmic drugs in the treatment of dry eye disease: A systematic literature review. *The ocular surface*. Jul 2019;17(3):412-423. doi:10.1016/j.jtos.2019.02.012
12. Milner MS, Beckman KA, Luchs JI, et al. Dysfunctional tear syndrome: dry eye disease and associated tear film disorders - new strategies for diagnosis and treatment. *Current opinion in ophthalmology*. Jan 2017;27 Suppl 1(Suppl 1):3-47. doi:10.1097/01.icu.0000512373.81749.b7

## Reimbursement Policy:

### Evaluation of Dry Eyes - Lab Benefit Program (LBM)

13. Tomlinson A, Khanal S, Ramaesh K, Diaper C, McFadyen A. Tear film osmolarity: determination of a referent for dry eye diagnosis. *Investigative ophthalmology & visual science*. Oct 2006;47(10):4309-15. doi:10.1167/iovs.05-1504
14. Akpek EK, Amescua G, Farid M, et al. Dry Eye Syndrome Preferred Practice Pattern. *Ophthalmology*. 2019;doi:10.1016/j.ophtha.2018.10.023
15. Baenninger PB, Voegeli S, Bachmann LM, et al. Variability of Tear Osmolarity Measurements With a Point-of-Care System in Healthy Subjects-Systematic Review. *Cornea*. Jul 2018;37(7):938-945. doi:10.1097/ico.0000000000001562
16. Honda N, Miyai T, Nejima R, et al. Effect of latanoprost on the expression of matrix metalloproteinases and tissue inhibitor of metalloproteinase 1 on the ocular surface. *Archives of ophthalmology (Chicago, Ill : 1960)*. Apr 2010;128(4):466-71. doi:10.1001/archophthalmol.2010.40
17. Chotikavanich S, de Paiva CS, Li de Q, et al. Production and activity of matrix metalloproteinase-9 on the ocular surface increase in dysfunctional tear syndrome. *Investigative ophthalmology & visual science*. Jul 2009;50(7):3203-9. doi:10.1167/iovs.08-2476
18. Sambursky R, Davitt WF, 3rd, Latkany R, et al. Sensitivity and specificity of a point-of-care matrix metalloproteinase 9 immunoassay for diagnosing inflammation related to dry eye. *JAMA ophthalmology*. Jan 2013;131(1):24-8. doi:10.1001/jamaophthalmol.2013.561
19. Messmer EM, von Lindenfels V, Garbe A, Kampik A. Matrix Metalloproteinase 9 Testing in Dry Eye Disease Using a Commercially Available Point-of-Care Immunoassay. *Ophthalmology*. Nov 2016;123(11):2300-2308. doi:10.1016/j.ophtha.2016.07.028
20. AXIM. AXIM EYE. Accessed 1/12/2023, [https://aximbiotech.com/wp-content/uploads/2021/11/Axim\\_Eye\\_Brochure\\_11-15-21\\_v3.pdf](https://aximbiotech.com/wp-content/uploads/2021/11/Axim_Eye_Brochure_11-15-21_v3.pdf)
21. Kanellopoulos AJ, Asimellis G. In pursuit of objective dry eye screening clinical techniques. *Eye and vision (London, England)*. 2016;3:1. doi:10.1186/s40662-015-0032-4
22. Brissette AR, Drinkwater OJ, Bohm KJ, Starr CE. The utility of a normal tear osmolarity test in patients presenting with dry eye disease like symptoms: A prospective analysis. *Cont Lens Anterior Eye*. Apr 2019;42(2):185-189. doi:10.1016/j.clae.2018.09.002
23. Tashbayev B, Utheim TP, Utheim ØA, et al. Utility of Tear Osmolarity Measurement in Diagnosis of Dry Eye Disease. *Scientific Reports*. 2020/03/26 2020;10(1):5542. doi:10.1038/s41598-020-62583-x
24. Chan TC, Ye C, Chan KP, Chu KO, Jhanji V. Evaluation of point-of-care test for elevated tear matrix metalloproteinase 9 in post-LASIK dry eyes. *Br J Ophthalmol*. Sep 2016;100(9):1188-91. doi:10.1136/bjophthalmol-2015-307607
25. Jun JH LY, Son MJ, Kim H Importance of tear volume for positivity of tear matrix metalloproteinase-9 immunoassay. *PLoS ONE*. 2020;15(7)doi:10.1371/journal.pone.0235408
26. Lee YH, Bang S-P, Shim K-Y, Son M-J, Kim H, Jun JH. Association of tear matrix metalloproteinase 9 immunoassay with signs and symptoms of dry eye disease: A cross-sectional study using qualitative, semiquantitative, and quantitative strategies. *PLoS one*. 2021;16(10):e0258203-e0258203. doi:10.1371/journal.pone.0258203
27. Choi M, Park YM, Ko BY. Comparative Evaluation of Matrix Metalloproteinase-9 Immunoassay and Tear Osmolarity Measurement for Diagnosing Severity of Dry Eye Disease. *Korean J Ophthalmol*. Oct 2023;37(5):409-416. doi:10.3341/kjo.2023.0062
28. Kawashima M, Kawakita T, Inaba T, et al. Dietary lactoferrin alleviates age-related lacrimal gland dysfunction in mice. *PLoS One*. 2012;7(3):e33148. doi:10.1371/journal.pone.0033148
29. Ponzini E, Scotti L, Grandori R, Tavazzi S, Zambon A. Lactoferrin Concentration in Human Tears and Ocular Diseases: A Meta-Analysis. *Investigative ophthalmology & visual science*. Oct 1 2020;61(12):9. doi:10.1167/iovs.61.12.9
30. AAO. Dry Eye Syndrome PPP 2023. <https://www.aaopt.org/education/preferred-practice-pattern/dry-eye-syndrome-ppp-2023>
31. AAO. Cornea/External Disease Summary Benchmarks - 2022. Updated December. <https://www.aaopt.org/education/summary-benchmark-detail/cornea-external-disease-summary-benchmarks-2020>

## Reimbursement Policy:

### Evaluation of Dry Eyes - Lab Benefit Program (LBM)

32. Craig JP, Nelson JD, Azar DT, et al. TFOS DEWS II Report Executive Summary. *The ocular surface*. Oct 2017;15(4):802-812. doi:10.1016/j.jtos.2017.08.003
33. AOA. Care of the Patient with Ocular Surface Disorders. <https://www.aoa.org/AOA/Documents/Practice%20Management/Clinical%20Guidelines/Consensus-based%20guidelines/Care%20of%20Patient%20with%20Ocular%20Surface%20Disorders.pdf>
34. Foulks GN, Forstot SL, Donshik PC, et al. Clinical guidelines for management of dry eye associated with Sjögren disease. *The ocular surface*. Apr 2015;13(2):118-32. doi:10.1016/j.jtos.2014.12.001
35. Starr CE, Gupta PK, Farid M, et al. An algorithm for the preoperative diagnosis and treatment of ocular surface disorders. *J Cataract Refract Surg*. May 2019;45(5):669-684. doi:10.1016/j.jcrs.2019.03.023
36. FDA. K934473. Updated 02/03/2025. Accessed 11/20/2018, 2018. <https://www.accessdata.fda.gov/scripts/cdrh/devicesatfda/index.cfm?db=pmn&id=K934473>
37. FDA. OcuSense, Inc., TearLab Osmolarity System [https://www.accessdata.fda.gov/cdrh\\_docs/pdf8/K083184.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf8/K083184.pdf)
38. Turker Medical. TearLab Osmolarity System. <https://trukera.com/>
39. FDA. 501(k) Summary of Safety and Effectiveness. [https://www.accessdata.fda.gov/cdrh\\_docs/pdf13/K132066.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf13/K132066.pdf)

## Revision History

Company(ies)	DATE	REVISION
<b>EmblemHealth</b>	4/2025	<ul style="list-style-type: none"> <li>Updated for clarity; no changes to coding or coverage criteria</li> </ul>
<b>EmblemHealth</b>	4/2025	<ul style="list-style-type: none"> <li>Transferred policy content to individual company-branded template. No changes to policy title or policy number.</li> </ul>
<b>EmblemHealth</b>	6/2024	<ul style="list-style-type: none"> <li>Lab Benefit Program (LBM) expanded to include EmblemHealth HMO/ PPO (Non-City) Commercial, Medicare and Medicaid plans effective 10/1/2024</li> </ul>
<b>EmblemHealth ConnectiCare</b>	5/2024	<ul style="list-style-type: none"> <li>Updated for clarity; no changes to coding or coverage criteria</li> </ul>
<b>EmblemHealth ConnectiCare</b>	7/2023	<ul style="list-style-type: none"> <li>Policy updated for clarity</li> </ul>
<b>EmblemHealth ConnectiCare</b>	11/2022	<ul style="list-style-type: none"> <li>Reformatted and reorganized policy, transferred content to new template with new Reimbursement Policy Number</li> </ul>