



Policy:	Ophthalmology – Vascular Endothelial Growth Factor Inhibitors – Beovu Utilization Management Medical Policy • Beovu <sup>®</sup> (brolucizumab intravitreal injection – Novartis)		
Date:		11/15/2023	
Applicable Lines of Business:		Medicare Advantage - Medical	
Applicable States:		NGS, J6: Wisconsin, Minnesota, Illinois NGS, JK: New York, Connecticut, Massachusetts, Maine, New Hampshire, Rhode Island, Vermont	

## Medicare Advantage Medical Utilization Review Policy

## **OVERVIEW**

Beovu, a vascular endothelial growth factor (VEGF) inhibitor, is indicated for the following uses:<sup>1</sup>

- Diabetic macular edema (DME).
- Neovascular (wet) age-related macular degeneration (nAMD).

For DME, the recommended dose for Beovu is 6 mg administered by intravitreal (IVT) injection every six weeks (every 39 to 45 days) for the first 5 doses, followed by 6 mg IVT injection once every 8 to 12 weeks. For nAMD, the recommended dose for Beovu is 6 mg administered by IVT injection every month (every 25 to 31 days) for the first 3 doses, followed by 6 mg IVT injection once every 8 to 12 weeks.

## **Other Uses with Supportive Evidence**

Overproduction of VEGF may lead to other eye conditions, including neovascular glaucoma, retinopathy of prematurity, and other retinal and choroidal neovascular conditions affecting the eye, the VEGF inhibitors also have the potential to be used off-label and reduce vision loss associated with other eye conditions related to increased VEGF production.<sup>2,3</sup> The use of anti-VEGF agents have been shown to stop the angiogenic process and maintain visual acuity and improve vision in patients with certain neovascular ophthalmic conditions; therefore, research is rapidly evolving on the use of VEGF inhibitors in other neovascular ophthalmic conditions which threaten vision.<sup>4,5</sup> Anti-VEGF therapy has the potential to be used off-label in other neovascular conditions affecting the eye and may prevent or slow visual impairment.<sup>2,4,5</sup>

#### **POLICY STATEMENT**

Prior authorization is recommended for medical benefit coverage of Beovu. Approval is recommended for those who meet the Criteria and Dosing for the listed indication(s). All approvals are provided for the duration noted below.

This policy incorporates Medicare coverage guidance as set forth in National Coverage Determinations (NCDs) and Local Coverage Determinations (LCDs), as well as in companion policy articles and other guidance applicable to the relevant service areas. These documents are cited in the References section of this policy. In some cases, this guidance includes specific lists of HCPCS and ICD-10 codes to help inform the coverage determination process. The Articles that include specific lists for billing and coding purposes will be included in the Reference section of this policy. However, to the extent that this policy cites such lists of HCPCS and ICD-10 codes, they should be used for reference purposes only. The presence of a specific HCPCS or ICD-10 code in a chart or companion article to an LCD is not by itself sufficient to approve coverage. Similarly, the absence of such a code does <u>not</u> necessarily mean that the applicable condition or diagnosis is excluded from coverage.

Ophthalmology – Vascular Endothelial Growth Factor Inhibitors – Beovu MA Medical UM Policy Page 2  $\,$ 

<u>Note</u>: Conditions for coverage outlined in this Medicare Advantage Medical Policy may be less restrictive than those found in applicable National Coverage Determinations, Local Coverage Determinations and/or Local Coverage Articles. Examples of situations where this clinical policy may be less restrictive include, but are not limited to, coverage of additional indications supported by CMS-approved compendia and the exclusion from this policy of additional coverage criteria requirements outlined in applicable National Coverage Determinations, Local Coverage Determinations and/or Local Coverage Determinations, Local Coverage Criteria requirements outlined in applicable National Coverage Determinations, Local Coverage Determinations and/or Local Coverage Articles.

# **RECOMMENDED AUTHORIZATION CRITERIA**

Coverage of Beovu is recommended in those who meet the following criteria.

## **FDA-Approved Indications**

1. Diabetic Macular Edema.

Criteria. Approve for 1 year.

**Dosing.** Approve if the requested dosing meets the following (A and B):

- A) The dose is 6 mg administered by intravitreal injection for each eye being treated; AND
- **B)** The dosing interval is not more frequent than once every 39 days for five doses, followed by not more frequently than once every 8 weeks for each eye being treated.

### 2. Neovascular (Wet) Age-Related Macular Degeneration.

Criteria. Approve for 1 year.

**Dosing.** Approve if the requested dosing meets the following (A and B):

- A) The dose is 6 mg administered by intravitreal injection for each eye being treated; AND
- **B)** The dosing interval is not more frequent than once every 25 days for three doses, followed by not more frequently than once every 8 weeks for each eye being treated.

#### **Other Uses with Supportive Evidence**

# 3. Other Neovascular Diseases of the Eye.

<u>Note</u>: Examples of other neovascular diseases of the eye include neovascular glaucoma, retinopathy of prematurity, sickle cell neovascularization, choroidal neovascular conditions.

Criteria. Approve for 1 year.

**Dosing.** Approve if the requested dosing meets the following (A <u>and</u> B):

- A) The dose is 6 mg administered by intravitreal injection for each eye being treated; AND
- **B)** The dosing interval is not more frequent than once every 25 days for three doses, followed by not more frequently than once every 8 weeks for each eye being treated.





11/15/2023 © 2023. All Rights Reserved.

## **CONDITIONS NOT RECOMMENDED FOR APPROVAL**

Coverage of Beovu is not recommended in the following situations:

**1.** Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

#### REFERENCES

- 1. Beovu® [prescribing information]. Hanover, NJ: Novartis; May 2022.
- 2. Barakat MR, Kaiser PK. VEGF inhibitors for the treatment of neovascular age-related macular degeneration. *Expert Opin Investig Drugs*. 2009;18(5):637-646.
- 3. Tolentino M. Systemic and ocular safety of intravitreal anti-VEGF therapies for ocular neovascular disease. *Surv Ophthalmol.* 2011;56(2):95-113.
- 4. Kinnunen K, Ylä-Herttuala S. Vascular endothelial growth factors in retinal and choroidal neovascular diseases. *Ann Med.* 2012;44(1):1-17.
- 5. Horsley MB, Kahook MY. Anti-VEGF therapy for glaucoma. Curr Opin Ophthalmol. 2010;21(2):112-117.
- Centers for Medicare and Medicaid Services, National Government Services, Inc, Local Coverage Determination (LCD): Drugs and Biologicals, Coverage of, for Label and Off-Label Uses (L33394) [original date 10/01/2015; revision effective date 11/1/2022]. Accessed on December 12, 2023.
- Centers for Medicare and Medicaid Services, National Government Services, Inc, Local Coverage Article (LCA): Billing and Coding: Ranibizumab, Aflibercept and Brolucizumab-dbll (A52451) [original date 10/01/2015; revision effective date 04/1/2023]. Accessed on December 12, 2023.

# HISTORY

Type of Revision	Summary of Changes*	Date
Policy created	New Medicare Advantage Medical Policy	02/05/2020
Policy revision	*Added the following to the Policy Statement " <u>Note</u> : Conditions for coverage outlined in this Medicare Advantage Medical Policy may be less restrictive than those found in applicable National Coverage Determinations, Local Coverage Determinations and/or Local Coverage Articles. Examples of situations where this clinical policy may be less restrictive include, but are not limited to, coverage of additional indications supported by CMS-approved compendia and the exclusion from this policy of additional coverage criteria requirements outlined in applicable National Coverage Determinations, Local Coverage Determinations and/or Local Coverage Articles."	08/07/2020
Policy revision	<b>Neovascular (Wet) Age-Related Macular Degeneration:</b> To align with the FDA-approved dosing, the dose was changed from " $\leq 6$ mg" to "is 6 mg". <b>Other Neovascular Diseases of the Eye:</b> Examples of other neovascular diseases of the eye were moved to a Note. To align with the FDA-approved dosing, the dose was changed from " $\leq 6$ mg" to "is 6 mg".	12/20/2021
Policy revision	Diabetic Macular Edema: This indication was added to the policy.	06/29/2022
Policy review	No criteria changes.	11/16/2022
Policy review	No criteria changes.	11/15/2023



© 2023. All Rights Reserved.