

# **Medical Policy:**

## Osteochondral Grafting

POLICY NUMBER	LAST REVIEW
MG.MM.SU.68bC	February 9, 2024

#### Medical Guideline Disclaimer Property of EmblemHealth. All rights reserved.

The treating physician or primary care provider must submit to EmblemHealth, or ConnectiCare, as applicable (hereinafter jointly referred to as "EmblemHealth"), the clinical evidence that the member meets the criteria for the treatment or surgical procedure. Without this documentation and information, EmblemHealth will not be able to properly review the request preauthorization or post-payment review. The clinical review criteria expressed below reflects how EmblemHealth determines whether certain services or supplies are medically necessary. This clinical policy is not intended to pre-empt the judgment of the reviewing medical director or dictate to health care providers how to practice medicine. Health care providers are expected to exercise their medical judgment in rendering appropriate care. Health care providers are expected to exercise their medical judgment in rendering appropriate

EmblemHealth established the clinical review criteria based upon a review of currently available clinical information (including clinical outcome studies in the peer reviewed published medical literature, regulatory status of the technology, evidence-based guidelines of public health and health research agencies, evidence-based guidelines and positions of leading national health professional organizations, views of physicians practicing in relevant clinical areas, and other relevant factors). EmblemHealth expressly reserves the right to revise these conclusions as clinical information changes and welcomes further relevant information. Each benefit program defines which services are covered. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that this service or supply is covered and/or paid for by EmblemHealth, as some programs exclude coverage for services or supplies that EmblemHealth considers medically necessary.

If there is a discrepancy between this guideline and a member's benefits program, the benefits program will govern. Identification of selected brand names of devices, tests and procedures in a medical coverage policy is for reference only and is not an endorsement of any one device, test or procedure over another. In addition, coverage may be mandated by applicable legal requirements of a state, the Federal Government or the Centers for Medicare & Medicaid Services (CMS) for Medicare and Medicaid members. All coding and web site links are accurate at time of publication.

EmblemHealth may also use tools developed by third parties, such as the MCG™ Care Guidelines, to assist us in administering health benefits. The MCG™ Care Guidelines are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice. EmblemHealth Services Company, LLC, has adopted this policy in providing management, administrative and other services to EmblemHealth Plan, Inc., EmblemHealth Insurance Company, EmblemHealth Services Company, LLC, and Health Insurance Plan of Greater New York (HIP) related to health benefit plans offered by these entities. ConnectiCare, an EmblemHealth company, has also adopted this policy. All of the aforementioned entities are affiliated companies under common control of EmblemHealth Inc.

#### **Definitions**

Autografts	Grafts of bone and cartilage harvested from either the patient's non-weight bearing surfaces (or surfaces that bear less weight), which is then implanted in the defect. Autografting is typically used to repair smaller defects.
	The advantages of using autograft material include graft availability, the absence of possible disease transmission risk, and that the procedure is a single-stage procedure. Disadvantages include donor site morbidity and limited available graft volume. In addition, tissue may have to be harvested from two different donor sites in order to provide enough material for a large defect without compromising the donor site.
	The COR and COR2 systems (DePuy-Mitek) can be utilized for autograft or allograft transplantation.
Allografts	Grafts of bone and cartilage harvested from a cadaver joint (may be fresh or cryopreserved), which is then implanted in the defect. Allografting is typically used to repair larger defects of the knee (e.g., 10 cm <sup>2</sup> ).
	The use of allograft cartilage has the advantage of providing osteochondral segments that are must be implanted within 72 hours of donor death, may be difficult to obtain (due to scarcity) and may also entail a

	concern of disease transmission.	
	Application of osteochondral allografting is limited because cryopreserved allografts do not contain an acceptable level of cartilage viability, and cryopreservation may decrease the viability of the cartilage cells.	
	A minimally processed osteochondral allograft (e.g., Chondrofix® [Zimmer]) is composed of decellularized hyaline cartilage and cancellous bone, which can be used "off the shelf" with precut cylinders of 7–15 mm. Multiple cylinders may be used to fill larger defects. The Osteochondral Autograft Transfer System (OATS) involves use of larger plugs usually filling the entire defect with a single plug.	
	The COR and COR2 systems can be utilized for autograft or allograft transplantation.	
Mosaicplasty	A technique of creating an osteochondral autograft by harvesting and transplanting multiple small cylindrical osteochondral plugs from the less weight-bearing periphery of the patellofemoral area and inserting them into drilled tunnels in the defective section of cartilage. (E.g., Mosaicplasty System)	

#### Guideline

Osteochondral autografting and allografting have been investigated for full-thickness cartilage defects of weight bearing surfaces secondary to trauma or conditions such as osteochondritis dissecans. The goal of grafting is to re-establish the cartilage matrix with chondrocytes and support bone in order to improve joint function and decrease pain.

Osteochondral autografting and allografting with one or more cores of osteochondral tissue is considered medically necessary for the treatment of cartilaginous defects caused by acute or repetitive trauma in the knee or ankle when all of the following criteria are met:

- 1. Presence of disabling, localized knee or ankle pain of  $\geq$  6 months duration that is refractory to conservative treatment, as well as abrasion arthroplasty and/or microfracture techniques
- 2. Mature skeletal growth achieved, or not a suitable candidate, for total joint replacement
- 3. Cartilage defect ≥ 1cm<sup>2</sup>
- 4. Focal, full thickness isolated defect of the weight bearing surface of any of the following areas:
  - a. Talus
  - b. Medial or lateral femoral condyles
  - c. Trochlear region of the knee
  - d. Patellar chondral defect
- 5. Defect is unipolar
- 6. Lesion is largely contained with near-normal surrounding articular cartilage and articulating cartilage (grades 0, 1, 2)
- 7. In the knee, meniscus is intact or has stable partial tears. (Partial meniscectomy may be indicated and does not preclude osteochondral grafting)
- 8. Normal or near-normal joint space with no > 15% joint space narrowing
- 9. No presence of active infection
- 10. No inflammation or osteoarthritis in the joint
- 11. Joint is stable with normal alignment. (A procedure to correct alignment may be performed in combination with, or prior to, grafting)

- 12. Member is willing and able to comply with post-operative weight-bearing restrictions and rehabilitation
- 13. No history of cancer in the bones, cartilage, fat or muscle of the affected limb

# **Limitations/Exclusions**

Osteochondral grafting is not considered medically necessary for any of the following due to insufficient evidence of therapeutic value:

- 1. Use in joints other than the knee and ankle (e.g., shoulder, elbow, hip)
- 2. Use of minced articular cartilage (autologous or allogeneic) (e.g., Cartilage Autograft Implantation System [CAIS], BioCartilage® [Arthrex], DeNovo NT Graft [natural tissue graft; ISTO Technologies]
- 3. Decellularized osteochondral allograft plugs (e.g., Chondrofix)
- 4. Reduced allograft discs (e.g., ProChondrix, Cartiform)

### **Procedure Codes**

27415	Osteochondral allograft, knee, open	
27416	Osteochondral autograft(s), knee, open (eg, mosaicplasty) (includes harvesting of autograft[s])	
28446	Open osteochondral autograft, talus (includes obtaining graft[s])	
29866	Arthroscopy, knee, surgical; osteochondral autograft(s) (eg, mosaicplasty) (includes harvesting of the autograft[s])	
29867	Arthroscopy, knee, surgical; osteochondral allograft (eg, mosaicplasty)	

## **Diagnosis Codes**

M22.40	Chondromalacia patellae, unspecified knee
M22.41	Chondromalacia patellae, right knee
M22.42	Chondromalacia patellae, left knee
M23.000	Cystic meniscus, unspecified lateral meniscus, right knee
M23.001	Cystic meniscus, unspecified lateral meniscus, left knee
M23.002	Cystic meniscus, unspecified lateral meniscus, unspecified knee
M23.003	Cystic meniscus, unspecified medial meniscus, right knee
M23.004	Cystic meniscus, unspecified medial meniscus, left knee
M23.005	Cystic meniscus, unspecified medial meniscus, unspecified knee
M23.006	Cystic meniscus, unspecified meniscus, right knee
M23.007	Cystic meniscus, unspecified meniscus, left knee
M23.009	Cystic meniscus, unspecified meniscus, unspecified knee
M23.011	Cystic meniscus, anterior horn of medial meniscus, right knee
M23.012	Cystic meniscus, anterior horn of medial meniscus, left knee

M23.019	Cystic meniscus, anterior horn of medial meniscus, unspecified knee	
M23.021	Cystic meniscus, posterior horn of medial meniscus, right knee	
M23.022	Cystic meniscus, posterior horn of medial meniscus, left knee	
M23.029	Cystic meniscus, posterior horn of medial meniscus, unspecified knee	
M23.031	Cystic meniscus, other medial meniscus, right knee	
M23.032	Cystic meniscus, other medial meniscus, left knee	
M23.039	Cystic meniscus, other medial meniscus, unspecified knee	
M23.041	Cystic meniscus, anterior horn of lateral meniscus, right knee	
M23.042	Cystic meniscus, anterior horn of lateral meniscus, left knee	
M23.049	Cystic meniscus, anterior horn of lateral meniscus, unspecified knee	
M23.051	Cystic meniscus, posterior horn of lateral meniscus, right knee	
M23.052	Cystic meniscus, posterior horn of lateral meniscus, left knee	
M23.059	Cystic meniscus, posterior horn of lateral meniscus, unspecified knee	
M23.061	Cystic meniscus, other lateral meniscus, right knee	
M23.062	Cystic meniscus, other lateral meniscus, left knee	
M23.069	Cystic meniscus, other lateral meniscus, unspecified knee	
M23.2	Derangement of meniscus due to old tear or injury	
M23.200	Derangement of unspecified lateral meniscus due to old tear or injury, right knee	
M23.201	Derangement of unspecified lateral meniscus due to old tear or injury, left knee	
M23.202	Derangement of unspecified lateral meniscus due to old tear or injury, unspecified knee	
M23.203	Derangement of unspecified medial meniscus due to old tear or injury, right knee	
M23.204	Derangement of unspecified medial meniscus due to old tear or injury, left knee	
M23.205	Derangement of unspecified medial meniscus due to old tear or injury, unspecified knee	
M23.206	Derangement of unspecified meniscus due to old tear or injury, right knee	
M23.207	Derangement of unspecified meniscus due to old tear or injury, left knee	
M23.209	Derangement of unspecified meniscus due to old tear or injury, unspecified knee	
M23.211	Derangement of anterior horn of medial meniscus due to old tear or injury, right knee	
M23.212	Derangement of anterior horn of medial meniscus due to old tear or injury, left knee	
M23.219	Derangement of anterior horn of medial meniscus due to old tear or injury, unspecified knee	
M23.221	Derangement of posterior horn of medial meniscus due to old tear or injury, right knee	
M23.222	Derangement of posterior horn of medial meniscus due to old tear or injury, left knee	
M23.229	Derangement of posterior horn of medial meniscus due to old tear or injury, unspecified knee	
M23.231	Derangement of other medial meniscus due to old tear or injury, right knee	
M23.232	Derangement of other medial meniscus due to old tear or injury, left knee	
M23.239	Derangement of other medial meniscus due to old tear or injury, unspecified knee	

M23.241	Derangement of anterior horn of lateral meniscus due to old tear or injury, right knee	
M23.242	Derangement of anterior horn of lateral meniscus due to old tear or injury, left knee	
M23.249	Derangement of anterior horn of lateral meniscus due to old tear or injury, unspecified knee	
M23.251	Derangement of posterior horn of lateral meniscus due to old tear or injury, right knee	
M23.252	Derangement of posterior horn of lateral meniscus due to old tear or injury, left knee	
M23.259	Derangement of posterior horn of lateral meniscus due to old tear or injury, unspecified knee	
M23.261	Derangement of other lateral meniscus due to old tear or injury, right knee	
M23.262	Derangement of other lateral meniscus due to old tear or injury, left knee	
M23.269	Derangement of other lateral meniscus due to old tear or injury, unspecified knee	
M23.300	Other meniscus derangements, unspecified lateral meniscus, right knee	
M23.301	Other meniscus derangements, unspecified lateral meniscus, left knee	
M23.302	Other meniscus derangements, unspecified lateral meniscus, unspecified knee	
M23.303	Other meniscus derangements, unspecified medial meniscus, right knee	
M23.304	Other meniscus derangements, unspecified medial meniscus, left knee	
M23.305	Other meniscus derangements, unspecified medial meniscus, unspecified knee	
M23.306	Other meniscus derangements, unspecified meniscus, right knee	
M23.307	Other meniscus derangements, unspecified meniscus, left knee	
M23.309	Other meniscus derangements, unspecified meniscus, unspecified knee	
M23.311	Other meniscus derangements, anterior horn of medial meniscus, right knee	
M23.312	Other meniscus derangements, anterior horn of medial meniscus, left knee	
M23.319	Other meniscus derangements, anterior horn of medial meniscus, unspecified knee	
M23.321	Other meniscus derangements, posterior horn of medial meniscus, right knee	
M23.322	Other meniscus derangements, posterior horn of medial meniscus, left knee	
M23.329	Other meniscus derangements, posterior horn of medial meniscus, unspecified knee	
M23.331	Other meniscus derangements, other medial meniscus, right knee	
M23.332	Other meniscus derangements, other medial meniscus, left knee	
M23.339	Other meniscus derangements, other medial meniscus, unspecified knee	
M23.341	Other meniscus derangements, anterior horn of lateral meniscus, right knee	
M23.342	Other meniscus derangements, anterior horn of lateral meniscus, left knee	
M23.349	Other meniscus derangements, anterior horn of lateral meniscus, unspecified knee	
M23.351	Other meniscus derangements, posterior horn of lateral meniscus, right knee	
M23.352	Other meniscus derangements, posterior horn of lateral meniscus, left knee	
M23.359	Other meniscus derangements, posterior horn of lateral meniscus, unspecified knee	
M23.361	Other meniscus derangements, other lateral meniscus, right knee	
M23.362	Other meniscus derangements, other lateral meniscus, left knee	
-		

M23.369	Other meniscus derangements, other lateral meniscus, unspecified knee	
M23.40	Loose body in knee, unspecified knee	
M23.41	Loose body in knee, right knee	
M23.42	Loose body in knee, left knee	
M23.50	Chronic instability of knee, unspecified knee	
M23.51	Chronic instability of knee, right knee	
M23.52	Chronic instability of knee, left knee	
M23.601	Other spontaneous disruption of unspecified ligament of right knee	
M23.602	Other spontaneous disruption of unspecified ligament of left knee	
M23.609	Other spontaneous disruption of unspecified ligament of unspecified knee	
M23.611	Other spontaneous disruption of anterior cruciate ligament of right knee	
M23.612	Other spontaneous disruption of anterior cruciate ligament of left knee	
M23.619	Other spontaneous disruption of anterior cruciate ligament of unspecified knee	
M23.621	Other spontaneous disruption of posterior cruciate ligament of right knee	
M23.622	Other spontaneous disruption of posterior cruciate ligament of left knee	
M23.629	Other spontaneous disruption of posterior cruciate ligament of unspecified knee	
M23.631	Other spontaneous disruption of medial collateral ligament of right knee	
M23.632	Other spontaneous disruption of medial collateral ligament of left knee	
M23.639	Other spontaneous disruption of medial collateral ligament of unspecified knee	
M23.641	Other spontaneous disruption of lateral collateral ligament of right knee	
M23.642	Other spontaneous disruption of lateral collateral ligament of left knee	
M23.649	Other spontaneous disruption of lateral collateral ligament of unspecified knee	
M23.671	Other spontaneous disruption of capsular ligament of right knee	
M23.672	Other spontaneous disruption of capsular ligament of left knee	
M23.679	Other spontaneous disruption of capsular ligament of unspecified knee	
M23.8X1	Other internal derangements of right knee	
M23.8X2	Other internal derangements of left knee	
M23.8X9	Other internal derangements of unspecified knee	
M23.90	Unspecified internal derangement of unspecified knee	
M23.91	Unspecified internal derangement of right knee	
M23.92	Unspecified internal derangement of left knee	
M25.861	Other specified joint disorders, right knee	
M25.862	Other specified joint disorders, left knee	
M25.869	Other specified joint disorders, unspecified knee	
M25.871	Other specified joint disorders, right ankle and foot	

M25.872	Other specified joint disorders, left ankle and foot	
M25.879	Other specified joint disorders, unspecified ankle and foot	
M92.40	Juvenile osteochondrosis of patella, unspecified knee	
M92.41	Juvenile osteochondrosis of patella, right knee	
M92.42	Juvenile osteochondrosis of patella, left knee	
M92.511	Juvenile osteochondrosis of proximal tibia, right leg	
M92.512	Juvenile osteochondrosis of proximal tibia, left leg	
M92.513	Juvenile osteochondrosis of proximal tibia, bilateral	
M92.519	Juvenile osteochondrosis of proximal tibia, unspecified leg	
M92.521	Juvenile osteochondrosis of tibia tubercle, right leg	
M92.522	Juvenile osteochondrosis of tibia tubercle, left leg	
M92.523	Juvenile osteochondrosis of tibia tubercle, bilateral	
M92.529	Juvenile osteochondrosis of tibia tubercle, unspecified leg	
M92.8	Other specified juvenile osteochondrosis	
M92.591	Other juvenile osteochondrosis of tibia and fibula, right leg	
M92.592	Other juvenile osteochondrosis of tibia and fibula, left leg	
M92.593	Other juvenile osteochondrosis of tibia and fibula, bilateral	
M92.599	Other juvenile osteochondrosis of tibia and fibula, unspecified leg	
M93.261	Osteochondritis dissecans, right knee	
M93.262	Osteochondritis dissecans, left knee	
M93.269	Osteochondritis dissecans, unspecified knee	
M93.271	Osteochondritis dissecans, right ankle and joints of right foot	
M93.272	Osteochondritis dissecans, left ankle and joints of left foot	
M93.279	Osteochondritis dissecans, unspecified ankle and joints of foot	
M94.261	Chondromalacia, right knee	
M94.262	Chondromalacia, left knee	
M94.269	Chondromalacia, unspecified knee	
M94.271	Chondromalacia, right ankle and joints of right foot	
M94.272	Chondromalacia, left ankle and joints of left foot	
M94.279	Chondromalacia, unspecified ankle and joints of foot	

# References

Astur DC, et al. Autologous osteochondral transplantation for treating patellar chondral injuries: evaluation, treatment, and outcomes of a two-year follow-up study. <u>J Bone Joint Surg Am</u> 2014 May 21;96(10):816-23.

Aubin PP, et al. Long-term follow-up of fresh femoral osteochondral allografts for posttraumatic knee defects. <u>Clin Orthopaed</u> Rel Res 2001;(2):S218-327.

Bekkers JE, et al. Treatment selection in articular cartilage lesions of the knee: a systematic review. Am J Sports Med 2009 Nov;37 Suppl 1:148S-55S.

Bekkers JE, et al. Articular cartilage evaluation after TruFit plug implantation analyzed by delayed gadolinium-enhanced MRI of cartilage (dGEMRIC). Am J Sports Med 2013 Jun;41(6):1290-5.

Bentley G, et al. Minimum ten-year results of a prospective randomized study of autologous chondrocyte implantation versus mosaicplasty for symptomatic articular cartilage lesions of the knee. J Bone Joint Surg Br 2012 Apr;94(4):504-9.

BlueCross BlueShield Association. Osteochondral autografts and allografts in the treatment of focal articular cartilage lesions. Medical Policy Reference Manual Policy #7.01.78. 2015 Jun 11.

Bugbee WD, et al. Bipolar fresh osteochondral allografting of the tibiotalar joint. <u>J Bone Joint Surg Am</u> 2013 Mar 6;95(5):426-32.

Camp CL, et al. Transplantation of a tibial osteochondral allograft to restore a large glenoid osteochondral defect. Orthopedics 2015 Feb;38(2):e147-52.

Chahal J, et al. Outcomes of osteochondral allograft transplantation in the knee. Arthroscopy 2013 Mar;29(3):575-88.

Chambers HG, et al. American Academy of orthopedic Surgeons clinical practice guideline on: the diagnosis and treatment of osteochondritis dissecans. J Bone Joint Surg Am 2012 Jul 18;94(14):1322-4.

De Caro F, et al. Large fresh osteochondral allografts of the knee: a systematic clinical and basic science review of the literature. Arthroscopy 2015 Apr;31(4):757-65.

Duif C, Koutah MA, Ackermann O, et al. Combination of autologous chondrocyte implantation (ACI) and osteochondral autograft transfer system (OATS) for surgical repair of larger cartilage defects of the knee joint. A review illustrated by a case report. Technol Health Care. 2015;23(5):531-537.

El-Rashidy H, et al. Fresh osteochondral allograft for the treatment of cartilage defects of the talus: a retrospective review. <u>J Bone Joint Surg Am</u> 2011 Sep 7;93(7):1634-40.

Emre TY, et al. Open mosaicplasty in osteochondral lesions of the talus: a prospective study. <u>J Foot Ankle Surg</u> 2012 Sep-Oct;51(5):556-60.

Erdil M, et al. Osteochondral autologous graft transfer system in the knee; mid-term results. Knee 2013 Jan;20(1):2-8.

Farr J, et al. Clinical, radiographic, and histological outcomes after cartilage repair with particulated juvenile articular cartilage: A 2-year prospective study. Am J Sports Med 2014 Jun;42(6):1417-25.

Figueroa D, et al. Osteochondral autografts in full thickness patella cartilage lesions. Knee 2011 Aug;18(4):220-3.

Galli MM, et al. Role of demineralized allograft subchondral bone in the treatment of shoulder lesions of the talus: clinical results with two-year follow-up. J Foot Ankle Surg 2014 Jul 9 [Epub ahead of print].

Gobbi A, et al. Osteochondral lesions of the talus: randomized controlled trial comparing chondroplasty, microfracture, and osteochondral autograft transplantation. <u>Arthroscopy</u> 2006 Oct;22(10):1085-92.

Gracitelli GC, et al. Fresh osteochondral allograft transplantation for isolated patellar cartilage injury. <u>Am J Sports Med</u> 2015 Apr;43(4):879-84.

Gracitelli GC, et al. Fresh osteochondral allografts in the knee: comparison of primary transplantation versus transplantation after failure of previous subchondral marrow stimulation. <u>Am J Sports Med</u> 2015 Apr;43(4):885-91.

Gudas R, et al. A prospective, randomized clinical study of osteochondral autologous transplantation versus microfracture for the treatment of osteochondritis dissecans in the knee joint in children. <u>J Pediatr Orthop</u> 2009 Oct-Nov;29(7):741-8.

Gudas R, et al. ten-year follow-up of a prospective, randomized clinical study of mosaic osteochondral autologous transplantation versus microfracture for the treatment of osteochondral defects in the knee joint of athletes. <u>Am J Sports Med</u> 2012 Nov;40(1):2499-508.

Gudas R, et al. Comparison of osteochondral autologous transplantation, microfracture, or debridement techniques in articular cartilage lesions associated with anterior cruciate ligament injury: a prospective study with a 3-year follow-up. <u>Arthroscopy</u> 2013 Jan;29(1):89-97.

Haasper C, et al. No mid-term difference in mosaicplasty in previously treated versus previously untreated patients with osteochondral lesions of the talus. <u>Arch Orthop Trauma Surg</u> 2008 May;128(5):499-504.

Haene R, et al. Intermediate outcomes of fresh talar osteochondral allografts for treatment of large osteochondral lesions of the talus. J Bone Joint Surg Am 2012 Jun 20;94(12):1105-10.

Hangody L, et al. Mosaicplasty for the treatment of articular defects of the knee and ankle. <u>Clin Orthopaed Rel Res</u> 2001 Oct;1(391):S328-36.

Hangody L, et al. Mosaicplasty for the treatment of osteochondritis dissecans of the talus: two to seven year results in 36 patients. <u>Foot Ankle Internat</u> 2001 Jul;22(7):552-8.

Hangody L, et al. Autologous osteochondral grafting- technique and long-term results. Injury 2008 Apr;39(Suppl)1:S32-9.

Harris JD, et al. Biological knee reconstruction: a systematic review of combined meniscal allograft transplantation and cartilage repair or restoration. <u>Arthroscopy</u> 2011;27(3):409-18.

Hettrich CM, et al. Cartilage repair: third-generation cell-based technologies- basic science, surgical techniques and clinical outcomes. <u>Sports Med Arthrosc</u> 2008 Dec;16(4):230-5.

Hindle P, et al. Autologous osteochondral mosaicplasty or TruFit<sup>™</sup> plugs for cartilage repair. <u>Knee Surg Sports Traumatol</u> Arthrosc 2014 Jun;22(6):1235-40.

Hu Y, et al. Treatment of large cystic medial osteochondral lesions of the talus with autologous osteoperiosteal cylinder grafts. Arthroscopy 2013 Aug;29(8):1372-9.

Imhoff AB, et al. Osteochondral transplantation of the talus: long-term clinical and magnetic resonance imaging evaluation. Am J Sports Med 2011 Jul;39(7):1487-93.

Iwasaki N, et al. Autologous osteochondral mosaicplasty for osteochondritis dissecans of the elbow in teenage athletes. <u>J Bone Joint Surg Am</u> 2009 Oct;91(10):2359-66.

Li Z, Zhu T, Fan W. Osteochondral autograft transplantation or autologous chondrocyte implantation for large cartilage defects of the knee: A meta-analysis. Cell Tissue Bank. 2016;17(1):59-67.Lynch TS, Patel RM, Benedick A, et al. Systematic review of autogenous osteochondral transplant outcomes. Arthroscopy. 2015;31(4):746-754.

Jeng CL, et al. Fresh osteochondral total ankle allograft transplantation for the treatment of ankle arthritis. <u>Foot Ankle Int</u> 2008 Jun;29(6):554-60.

Joshi N, et al. Synthetic resorbable scaffolds for the treatment of isolated patellofemoral cartilage defects in young patients: magnetic resonance imaging and clinical evaluation. <u>Am J Sports Med</u> 2012 Jun;40(6):1289-95.

Kim YS, et al. Clinical comparison of the osteochondral autograft transfer system and subchondral drilling in the osteochondral defects of the first metatarsal head. <u>Am J Sports Med</u> 2012 Aug;40(8):1824-33.

Kim YS, et al. Factors associated with the clinical outcomes of the osteochondral autograft system in osteochondral lesions of the talus: second-look arthroscopic evaluation. <u>Am J Sports Med</u> 2012 Dec;40(12):2709-19.

Kircher J, et al. Osteochondral autologous transplantation for the treatment of full-thickness cartilage defects of the shoulder: results at nine years. J Bone Joint Surg Br 2009 Apr;91(4):499-503.

Kon E, et al. How to treat osteochondritis dissecans of the knee: surgical techniques and new trends: AAOS exhibit selection. J Bone Joint Surg Am 2012 Jan 4;94(1):e1(1-8).

Kreuz P, et al. Mosaicplasty with autogenous talar autograft for osteochondral lesions of the talus after failed primary arthroscopic management – a prospective study with a 4-year follow-up. <u>Am J Sports Med</u> 2006:34(1):55-63.

Kyrch AJ, et al. Return to athletic activity after osteochondral allograft transplantation in the knee. <u>Am J Sports Med</u> 2012 May;40(5):1053-9.

Kyrch AJ, et al. Activity levels are higher after osteochondral autograft transfer mosaicplasty than after microfracture for articular

cartilage defects of the knee: a retrospective comparative study. J Bone Joint Surg Am 2012 Jun 6;94(11):971-8.

Lim HC, et al. Current treatments of isolated articular cartilage lesions of the knee achieve similar outcomes. <u>Clin Orthop Relat Res</u> 2012 Aug;470(8):2261-7.

Liu W, et al. Osteochondral autograft transplantation for acute osteochondral fractures associated with an ankle fracture. <u>Foot Ankle Int</u> 2011 Apr;32(4):437-42.

Lynch TS, et al. Systematic review of autogenous osteochondral transplant outcomes. Arthroscopy 2015 Apr;31(4):746-54.

Ma HL, et al. Osteochondral autografts transfer for post-traumatic osteochondral defect of the knee- 2-5 years follow-up. <u>Injury</u> 2004 Dec;35(12):1286-92.

Maruyama M, et al. Outcomes of an open autologous osteochondral plug graft for capitellar osteochondritis dissecans: time to return to sports. Am J Sports Med 2014 Sep;42(9):2122-77.

Meric G, et al. Fresh osteochondral allograft transplantation for bipolar reciprocal osteochondral lesions of the knee. Am <u>J Sports</u> Med 2015 Mar;43(3):709-14.

Murphy RT, et al. Osteochondral allograft transplantation of the knee in the pediatric and adolescent population. <u>Am J Sports Med</u> 2014 Mar;42(3):635-40.

National Institute for Health and Clinical Excellence. Mosaicplasty for symptomatic articular cartilage defects of the knee. March 2018. https://www.nice.org.uk/guidance/ipg607. Accessed February 9, 2024.

Ollat D, et al. Mosaic osteochondral transplantations in the knee joint, midterm results of the SFA multicenter study. <u>Orthop</u> Traumatol Surg Res 2011 Dec;97(8 Suppl):S160-6.

Ovesen J, et al. The clinical outcomes of mosaicplasty in the treatment of osteochondritis dissecans of the distal humeral capitellum of young athletes. J Shoulder Elbow Surg 2011 Jul;20(5):813-8.

Pearce CJ, et al. Synthetic osteochondral grafting of ankle osteochondral lesions. Foot Ankle Surg 2012 Jun;18(2):114-8.

Peterson W, et al. Osteochondral transplantation for the treatment of osteochondral defects at the talus with the Diamond twin system and graft harvesting from the posterior femoral condyles. <u>Arch Orthop Trauma Surg</u> 2014 Jun;134(6):843-52.

Raikin SM. Fresh osteochondral allografts for large-volume cystic osteochondral defects of the talus. <u>J Bone Joint Surg Am</u> 2009 Dec;91(12):2818-26.

Richter DL, Schenck RC Jr, Wascher DC, Treme G. Knee articular cartilage repair and restoration techniques: A review of the literature. Sports Health. 2016;8(2):153-160.

Sagstetter PJ, et al. Donor-site morbidity after osteochondral autologous transplantation for lesions of talus. <u>J Bone Joint Surg Am</u> 2009 Jul;91(17):1683-8.

Shimada K, et al. Cylindrical costal osteochondral autograft for reconstruction of large defects of the capitellum due to osteochondritis dissecans. J Bone Joint Surg Am 2012 Jun 6;94(11):992-1002.

Solheim E, et al. Results at 10 to 14 years after osteochondral autografting (mosaicplasty) in articular cartilage defects in the knee. Knee 2013 Aug;20(4):287-90.

Spak RT, et al. Fresh osteochondral allografts for patellofemoral arthritis – long-term followup. <u>Clin Ortho Rel Res</u> 2006 Mar;(444):193-200.

Ulstein S, et al. Microfracture technique versus osteochondral autologous transplantation mosaicplasty in patients with articular chondral lesions of the knee: a prospective randomized trial with long-term follow-up. Knee Surg Sports Traumatol Arthrosc 2014 Jun;22(6):1207-15.

Valderrabano V, et al. Reconstruction of osteochondral lesions of the talus with autologous spongiosa grafts and autologous matrix-induced chondrogenesis. <u>Am J Sports Med</u> 2013 Mar;41(3):519-27.

Vogt S, et al. Osteochondral transplantation in the elbow leads to good clinical and radiologic long-term results: An 8-to 14-year follow-up examination. <u>Am J Sports Med</u> 2011 Dec;39(12):2619-25.

Wang CJ. Treatment of focal articular cartilage lesions of the knee with autogenous osteochondral grafts – a 2 to 4 year follow-up

study. Arch Orthoped Trauma Surg 2002;122:169-72.

Woelfe JV, et al. Indications and limitations of osteochondral autologous transplantation in osteochondritis dissecans of the talus. Knee Surg Sports Traumatol Arthrosc 2013 Aug;21(8):1925-30.

Wu WC, et al. Autologous osteochondral graft for traumatic defects of finger joints. <u>J Hand Surg Eur Vol</u> 2012 Mar;37(3):251-7.

Yoon HS, et al. Osteochondral autologous transplantation is superior to repeat arthroscopy for the treatment of osteochondral lesions of the talus after failed primary arthroscopic treatment. <u>Am J Sports Med</u> 2014 Jun 6;42(8):18961903.

Solheim E, et al. Results at 10-14 years after microfracture treatment of articular cartilage defects in the knee. <u>Knee Surg Sports Traumatol Arthrosc</u> 2014 Nov 23 [Epub ahead of print].

Specialty matched clinical peer review.

### **Revision History**

Company(ies)	DATE	REVISION
EmblemHealth ConnectiCare	Feb. 10, 2023	Added patellar chondral defect indication to coincide with covered diagnostic coding listed in policy
ConnectiCare	Jan. 14, 2022	Retired MCG criteria for this service and adopted the clinical criteria of parent corporation EmblemHealth
EmblemHealth	Jan. 11, 2019	Added decellularized osteochondral allograft plugs (e.g., Chondrofix) and reduced allograft discs (e.g., ProChondrix, Cartiform) to Limitations/Exclusions section as investigational