

## Yescarta (axicabtagene ciloleucel)

Last Review Date: March 30, 2018  
Effective Date: April 1, 2018

Number: MG.MM.PH.42

### Medical Guideline Disclaimer

Property of EmblemHealth. All rights reserved. The treating physician or primary care provider must submit to EmblemHealth the clinical evidence that the patient meets the criteria for the treatment or surgical procedure. Without this documentation and information, EmblemHealth will not be able to properly review the request for prior authorization. The clinical review criteria expressed below reflects how EmblemHealth determines whether certain services or supplies are medically necessary. EmblemHealth established the clinical review criteria based upon a review of currently available clinical information (including clinical outcome studies in the peer-reviewed published medical literature, regulatory status of the technology, evidence-based guidelines of public health and health research agencies, evidence-based guidelines and positions of leading national health professional organizations, views of physicians practicing in relevant clinical areas, and other relevant factors). EmblemHealth expressly reserves the right to revise these conclusions as clinical information changes, and welcomes further relevant information. Each benefit program defines which services are covered. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that this service or supply is covered and/or paid for by EmblemHealth, as some programs exclude coverage for services or supplies that EmblemHealth considers medically necessary. If there is a discrepancy between this guideline and a member's benefits program, the benefits program will govern. In addition, coverage may be mandated by applicable legal requirements of a state, the Federal Government or the Centers for Medicare & Medicaid Services (CMS) for Medicare and Medicaid members. All coding and web site links are accurate at time of publication. EmblemHealth Services Company LLC, ("EmblemHealth") has adopted the herein policy in providing management, administrative and other services to HIP Health Plan of New York, HIP Insurance Company of New York, Group Health Incorporated and GHI HMO Select, related to health benefit plans offered by these entities. All of the aforementioned entities are affiliated companies under common control of EmblemHealth Inc.

### Definition

Yescarta (axicabtagene ciloleucel): is a CD19-directed genetically modified autologous T cell immunotherapy that kills CD19-expressing cancer cells. T cells from the patient are harvested and genetically modified ex vivo by retroviral transduction to express a chimeric antigen receptor (CAR), and after infusion back into the patient, results in CD28 and CD3-zeta co-stimulatory domains to activate causing T-cell activation, proliferation, acquisition of effector functions and secretion of inflammatory cytokines and chemokines.

Yescarta (axicabtagene ciloleucel) is FDA approved for the treatment of adult patients with relapsed or refractory large B-cell lymphoma after two or more lines of systemic therapy, including diffuse large B-cell lymphoma (DLBCL) not otherwise specified, primary mediastinal large B-cell lymphoma, high grade B-cell lymphoma, and DLBCL arising from follicular lymphoma.

Yescarta (axicabtagene ciloleucel) is available as cell suspension for infusion. Yescarta comprises a suspension of  $2 \times 10^6$  CAR-positive viable T cells per kg of body weight, with a maximum of  $2 \times 10^8$  CAR-positive viable T cells in approximately 68 mL.

### Guideline

Provider must submit documentation (which may include office notes and lab results) supporting that the patient has met all approval criteria.

Yescarta is considered medically necessary when all the below criteria are met:

- Patient is aged 18 years or greater; **AND**
- Patient has a confirmed diagnosis of large B-cell lymphoma, including:
  - Diffuse large B-cell lymphoma (DLBCL) not otherwise specified; **OR**
  - Primary mediastinal large B-cell lymphoma; **OR**
  - High grade B-cell lymphoma; **OR**
  - DLBCL arising from follicular lymphoma; **AND**

- The large B-cell lymphoma must be relapsed or refractory after two or more types of systemic therapy.
- Patient's disease is relapsed or refractory defined as one of the following:
  - Relapse within 1 year after autologous hematopoietic stem cell transplantation (HSCT);  
**OR**
  - Refractory disease to the most recent therapy; **AND**
- Patient did not receive prior allogeneic hematopoietic stem cell transplantation (HSCT); **AND**
- Patient has an ECOG performance status of 0-1; **AND**
- Patient has CD19-positive disease; **AND**
- Patient must not be currently pregnant and sexually-active females of reproductive potential should have pregnancy status verified through a pregnancy test; **AND**
- Patient does not have a clinically significant active systemic infection or inflammatory disorder; **AND**
- Patient has not received live vaccines within 6 weeks prior to the start of lymphodepleting chemotherapy, during Yescarta treatment, and will not receive live vaccines until immune recovery following treatment; **AND**
- Patient has been screened for hepatitis B virus (HBV), hepatitis C virus (HCV), and human immunodeficiency virus (HIV) in accordance with clinical guidelines prior to collection of cells (leukapheresis); **AND**
- Prophylaxis for infection has been followed according to local guidelines; **AND**
- Patient will be using Yescarta in conjunction with lymphodepleting chemotherapy (fludarabine 30 mg/m<sup>2</sup> daily for 3 days and cyclophosphamide 500 mg/m<sup>2</sup> daily on the fifth, fourth, and third day before infusion of Yescarta ; **AND**
- Healthcare facility has enrolled in the Yescarta REMS and training has been given to providers on the management of cytokine release syndrome (CRS) and neurological toxicities; **AND**
- Patient will be using Yescarta (axicabtagene cilileucel) at a treatment center that is certified to administer Yescarta (axicabtagene cilileucel); **AND**
- Patient will be monitored for signs and symptoms of Cytokine Release Syndrome (CRS) for at least 4 weeks after treatment with Yescarta (axicabtagene cilileucel) and will be counselled to seek immediate medical attention should signs and symptoms of CRS or a neurological event occur at any time; **AND**
- Patient will stay within proximity (within 2 hours) of the Yescarta (axicabtagene cilileucel) infusion center for at least 4 weeks following infusion.

### Limitations/Exclusions

- Dosing of Yescarta (axicabtagene cilileucel) is based on the number of chimeric antigen receptor (CAR)-positive viable T-cells.
- The target dose is : 2 X 10<sup>6</sup> CAR-positive viable T cells per kg body weight, with a maximum of 2 X 10<sup>8</sup> CAR- positive viable T cells.
- Patient has a diagnosis of primary central nervous system lymphoma
- Patient has previously received CAR-T therapy

### Initial Approval

- Approval will be granted for 1 single dose of Yescarta
- Coverage cannot be renewed, a maximum of one dose per lifetime will apply

### Applicable Procedure Codes

| HCPCS | HCPCS Description  |
|-------|--|
| Q2041 | Axicabtagene Ciloleucel, up to 200 million autologous Anti-CD19 CAR T Cells, including leukapheresis and dose preparation procedures, per infusion |

### Applicable ICD-10 Codes

| ICD-10 | ICD-10 Description  |
|--------|---|
| C82.20 | Follicular lymphoma grade III, unspecified, unspecified site                              |
| C82.21 | Follicular lymphoma grade III, unspecified, lymph nodes of head, face and neck            |
| C82.22 | Follicular lymphoma, grade III, unspecified, intrathoracic lymph nodes                    |
| C82.23 | Follicular lymphoma grade III, unspecified, intra-abdominal lymph nodes                   |
| C82.24 | Follicular lymphoma grade III, unspecified, lymph nodes of axilla and upper limb          |
| C82.25 | Follicular lymphoma grade III, unspecified, lymph nodes of inguinal region and lower limb |
| C82.26 | Follicular lymphoma grade III, unspecified, intrapelvic lymph nodes                       |
| C82.27 | Follicular lymphoma grade III, unspecified, spleen  |
| C82.28 | Follicular lymphoma grade III, unspecified, lymph nodes of multiple sites                 |
| C82.29 | Follicular lymphoma grade III, unspecified, extranodal and solid organ sites              |
| C82.30 | Follicular lymphoma grade IIIa, unspecified site  |
| C82.31 | Follicular lymphoma grade IIIa, lymph nodes of head, face and neck                        |
| C82.32 | Follicular lymphoma, grade IIIa, intrathoracic lymph nodes                                |
| C82.33 | Follicular lymphoma grade IIIa, intra-abdominal lymph nodes                               |
| C82.34 | Follicular lymphoma grade IIIa, lymph nodes of axilla and upper limb                      |
| C82.35 | Follicular lymphoma grade IIIa, lymph nodes of inguinal region and lower limb             |
| C82.36 | Follicular lymphoma grade IIIa, intrapelvic lymph nodes                                   |
| C82.37 | Follicular lymphoma grade IIIa, spleen  |
| C82.38 | Follicular lymphoma grade IIIa, lymph nodes of multiple sites                             |
| C82.39 | Follicular lymphoma grade IIIa, extranodal and solid organ sites                          |
| C82.40 | Follicular lymphoma grade IIIb, unspecified site  |
| C82.41 | Follicular lymphoma grade IIIb, lymph nodes of head, face and neck                        |
| C82.42 | Follicular lymphoma, grade IIIb, intrathoracic lymph nodes                                |
| C82.43 | Follicular lymphoma grade IIIb, intra-abdominal lymph nodes                               |
| C82.44 | Follicular lymphoma grade IIIb, lymph nodes of axilla and upper limb                      |
| C82.45 | Follicular lymphoma grade IIIb, lymph nodes of inguinal region and lower limb             |
| C82.46 | Follicular lymphoma grade IIIb, intrapelvic lymph nodes                                   |
| C82.47 | Follicular lymphoma grade IIIb, spleen  |

|        |   |
|--------|---|
| C82.48 | Follicular lymphoma grade IIIb, lymph nodes of multiple sites                               |
| C82.49 | Follicular lymphoma grade IIIb, extranodal and solid organ sites                            |
| C83.30 | Diffuse large B-cell lymphoma unspecified site  |
| C83.31 | Diffuse large B-cell lymphoma, lymph nodes of head, face, and neck                          |
| C83.32 | Diffuse large B-cell lymphoma intrathoracic lymph nodes                                     |
| C83.33 | Diffuse large B-cell lymphoma intra-abdominal lymph nodes                                   |
| C83.34 | Diffuse large B-cell lymphoma lymph nodes of axilla and upper limb                          |
| C83.35 | Diffuse large B-cell lymphoma, lymph nodes of inguinal region and lower limb                |
| C83.36 | Diffuse large B-cell lymphoma intrapelvic lymph nodes                                       |
| C83.37 | Diffuse large B-cell lymphoma, spleen   |
| C83.38 | Diffuse large B-cell lymphoma lymph nodes of multiple sites                                 |
| C83.39 | Diffuse large B-cell lymphoma extranodal and solid organ sites                              |
| C85.20 | Mediastinal (thymic) large B-cell lymphoma, unspecified site                                |
| C85.21 | Mediastinal (thymic) large B-cell lymphoma, lymph nodes of head, face and neck              |
| C85.22 | Mediastinal (thymic) large B-cell lymphoma, intrathoracic lymph nodes                       |
| C85.23 | Mediastinal (thymic) large B-cell lymphoma, intra-abdominal lymph nodes                     |
| C85.24 | Mediastinal (thymic) large B-cell lymphoma, lymph nodes of axilla and upper limb            |
| C85.25 | Mediastinal (thymic) large B-cell lymphoma, lymph nodes of inguinal region and lower limb   |
| C85.26 | Mediastinal (thymic) large B-cell lymphoma, intrapelvic lymph nodes                         |
| C85.27 | Mediastinal (thymic) large B-cell lymphoma, spleen  |
| C85.28 | Mediastinal (thymic) large B-cell lymphoma, lymph nodes of multiple sites                   |
| C85.29 | Mediastinal (thymic) large B-cell lymphoma, extranodal and solid organ sites                |
| C85.80 | Other specified types of non-Hodgkin lymphoma, unspecified site                             |
| C85.81 | Other specified types of non-Hodgkin lymphoma, lymph nodes of head, face and neck           |
| C85.82 | Other specified types of non-Hodgkin lymphoma, intrathoracic lymph nodes                    |
| C85.83 | Other specified types of non-Hodgkin lymphoma, intra-abdominal lymph nodes                  |
| C85.84 | Other specified types of non-Hodgkin lymphoma, lymph nodes of axilla and upper limb         |
| C85.85 | Other specified types of non-Hodgkin lymphoma, lymph nodes of inguinal region of lower limb |
| C85.86 | Other specified types of non-Hodgkin lymphoma, intrapelvic lymph nodes                      |
| C85.87 | Other specified types of non-Hodgkin lymphoma, spleen                                       |
| C85.88 | Other specified types of non-Hodgkin lymphoma, lymph nodes of multiple sites                |
| C85.89 | Other specified types of non-Hodgkin lymphoma, extranodal and solid organ sites             |
| Z85.72 | Personal history of non-Hodgkin lymphomas   |

## References

1. Yescarta [package insert]. Santa Monica, CA; Kite Pharma, Inc., October 2017.
2. Clinical Pharmacology Elsevier Gold Standard. 2017.
3. National Comprehensive Cancer Network. Cancer Guidelines and Drugs and Biologics Compendium. 2017.